

Alcohol Related Presentations to Lakes District Hospital's Emergency Department: January to December 2019

Prepared by the Public Health Service of Southern DHB September 2020

Key findings -

Annual summary 2019

- There were 607 alcohol related presentations to ED (5%)
- Three out of four months, an average of one person presented to ED with a life threatening alcohol related condition (triage level 1).
- Every week, an average of 5 people presented to ED with potentially life-threatening alcohol related conditions (triage levels 2 or 3).
- Alcohol related cases present in ED at any time of the day or night with the busiest times being Saturday and Sunday mornings (midnight-6am).
- April was the busiest month.
- 36% of cases were aged 25-34 years while 35% were aged between 18-24 years. Males presented more frequently than females (60% of presentations were male).

Background

From July 2017, all Emergency Departments (EDs) are mandated to record whether or not alcohol has been a factor in each admission. Alcohol presentation categories are shown below.

No - alcohol consumption is NOT directly associated with this presentation

Not known - couldn't determine if alcohol is associated with this presentation

Secondary - consequence of others' alcohol consumption

Yes - alcohol consumption is directly associated with this presentation

The presentation descriptions of interest are defined into two subgroups:

- Primary Principal person admitted who is affected by alcohol (coded above as Yes)
- Secondary Person who is admitted with injuries caused by an alcohol affected person

Unless otherwise specified, when this report refers to alcohol related presentations it includes both primary and secondary.

Limitations: Data quality is only as good as data entry; it is currently unknown if this process is consistent across all facilities who collect this data. This is an ongoing piece of work for Southern DHB.

Residents of Wanaka and surroundings may access facilities at Dunstan Hospital due to weather conditions closing the Crown Range. This report also does not capture transfer of ED admissions transferred through to Southland Hospital for subsequent treatment.

Triaging definition: Triaging is the process of deciding how serious a person's illness or injury is in order to prioritise cases so they can be treated by urgency. All presentations to ED are assigned a triage level, as below.

- 1 immediately life threatening
- 2 imminently life-threatening, or important time-critical (>10 min)
- **3** potentially life-threatening, potential adverse outcomes from delay > 30 min, or severe discomfort or distress
- **4** potentially serious, or potential adverse outcomes from delay > 60 min, or significant complexity or severity, or discomfort or distress
- 5 less urgent, or dealing with administrative issues only

Lakes District Hospital Emergency Data

This data reflects only acute harm that presented at the Lakes District Emergency Department. It does not include presentations to other parts of the health system or the full range of known harms from alcohol and their costs to both individuals and their communities. Alcohol consumption is a causal factor in more than 200 disease and injury conditions. Individual harm is also known to be related to the pattern of alcohol consumption: ²

- binge drinking can lead to interpersonal violence and injuries and result in emotional trauma;
- chronic medium to high level consumption is linked with liver and cardiovascular disease, mental health disorders and domestic violence;
- lower level consumption over long periods has been causally linked with a range of cancers.

The data provided is not representative of patterns and is not generalisable.

Table 1: ED presentations by alcohol classification (January-December 2019)

Alcohol Presentation description	Lakes District Hospital
No - alcohol consumption is NOT directly associated with this presentation	12343
Not known - couldn't determine if alcohol is associated with this presentation	102
Secondary - consequence of others' alcohol consumption	59
Yes - alcohol consumption is directly associated with this presentation	548
Grand Total	13052

There were 607 ED alcohol related presentations (5% of the total).

- Crudely 50 ED presentations every month, 1-2 every day
- 1 in 10 of these presentations were coded as secondary alcohol harm (10%).

Tables 2 and 3 show the profiles of the alcohol related ED presentations (n=607); age and gender as well as ethnicity.

Table 2: Alcohol related presentations were heavily skewed towards the younger age groups.

- 1. highest users were 25-34 year olds, closely followed by 18-24 year (almost three quarters of all presentations)
- 2. female presentations were typically less than males (40% v 60%)

Table 3: the majority of alcohol related presentations self-identified as European (83%), followed by Māori (7%)

Table 2: ED Alcohol related presentations by gender and age (January-December 2019)

		•		7 8					•	Grand
Age Range	< 15	15-17	18-24	25-34	35-44	45-54	55-64	65-74	75 +	Total
Female	2	9	98	79	32	7	12	3	1	243
Male	2	4	115	139	56	23	11	9	4	363
Undefined				1						1
Grand Total	4	13	213	219	88	30	23	12	5	607
Percentage	1%	2%	35%	36%	14%	5%	4%	2%	1%	100%

Table 3: ED Alcohol related presentations by ethnicity (January-December 2019)

Ethnicity	European	Māori	Pacific Peoples	Other Ethnicity	Asian	Grand Total
Total	505	42	10	21	21	599
Percentage	84%	7%	2%	4%	4%	100%

Note 1: there were 8 individuals (1%) where ethnicity was not defined

Note 2: some percentages do not total 100% due to rounding

Note 3: this data (both tables) should be compared against the latest Census data

¹ World Health Organization. Alcohol. Fact sheet; 2015, http://www.who.int/mediacentre/factsheets/fs349/en/

² Ministerial Council on Drug Strategy. Alcohol in Australia: issues and strategies; 2001

Successful and the second of t

Figure 1: ED alcohol related presentations by day and time (January-December 2019)

Note: Not appropriate to plot averages as numbers are too small

Alcohol related ED presentations are avoidable. The harm that people cause to themselves is *bad enough* but they also divert time and resources from other patients. The number of intoxicated people turning up in ED puts unnecessary strain on the service "people can be rude, aggressive, or in the worst circumstances — even violent towards doctors and nurses". This behaviour is not always limited to the individual requiring medical care as they may also be accompanied by associates who have also consumed high levels of alcohol. On these occasions, hospital security staff maybe required to restrain the patient and/or associate which is once again an avoidable drain on resources.


Figure 1 shows:

• Saturday and Sunday early mornings midnight-6am are disproportionately the busiest times

Figure 2 below shows:

- 4.7% of all ED presentations were alcohol related
- April had the highest proportion of alcohol related presentations (6.1%)

Figure 2: Proportion of ED presentations that are alcohol related (January-December 2019)



Note: Health cannot predict or comment on peaks and troughs as we cannot show a causal relationship to admission rates

Table 4: ED Alcohol related presentations: what does this harm look like?

ED Triage level	Primary alcohol harm	Secondary alcohol harm	Grand Total
1: Immediately life-threatening	9		9
2: Imminently life-threatening, or important time-critical (>10			
min)	49	6	55
3: Potentially life-threatening, potential adverse outcomes			
from delay > 30 min, or severe discomfort or distress	190	20	210
4: Potentially serious, or potential adverse outcomes from			
delay > 60 min, or significant complexity or severity, or			
discomfort or distress	277	28	305
5: Less urgent, or dealing with administrative issues only	23	5	28
Grand Total	548	59	607

Table 4 describes the burden of alcohol related presentations on ED by triage level:

- Every week, an average of 5 individuals presented who were in imminent or potentially lifethreatening conditions (triage levels 2-3)
- Three out of four months, an average of one person presented to ED with an immediately life threatening alcohol related condition (triage level 1)³

Table 5 describes the duration of stay for alcohol related presentations (n=607) in EDs.

- Almost 40% were discharged from ED within an hour
- 6% were in ED for longer than three hours but not admitted.
- Just over a quarter were subsequently admitted to a ward (ED or General) of which an unknown quantity (requiring surgery or medical interventions) were transferred to Southland Hospital, on occasions to Dunedin Public Hospital.

Table 5: ED Alcohol related presentations: level of harm measured

Duration of stay	Total	Percentage
ED only	438	72%
< 60 minutes	229	38%
1-2 hours	108	18%
2-3 hours	66	11%
> 3 hours	35	6%
Admission to ward	169	28%
Grand Total	607	100%

³ Triage levels 1-2 descriptions include: motor vehicle/scooter accident, self-harm (attempted suicides, overdoses), head injuries (assault, loss of consciousness), orthopaedic injuries, impalement of sharp objects/skin lacerations, confusion/semi-responsive, cardiovascular, alleged sexual assault