

# Management of Anaphylaxis

Acute onset of life-threatening airway and/or breathing and/or circulation problems and usually skin and/or mucosal changes.

## ASSESS

- Airway: swelling, hoarseness, noisy breathing (stridor)
- Breathing: fast, wheeze, cyanosis, fatigue, confusion
- Circulation: pale, clammy, slow capillary refill, low BP, faintness, drowsy/coma
- Skin and mucosal changes: urticaria, flushing, angioedema

**1. CALL FOR HELP** Send for emergency medical assistance (ambulance, doctor).

**2. POSITION PATIENT SAFELY** Do not allow them to stand and never leave them alone.



**3. ADMINISTER ADRENALINE** By deep IM injection into outer thigh.

Adrenaline dosage for 1:1,000 formulation is 0.01 mL/kg up to a maximum of 0.5 mL.

<b>AGE</b>	12 years and over	<b>DOSE</b>	500 mcg (0.5 mL)
------------	-------------------	-------------	------------------

Expect to see some response to the adrenaline within 1–2 minutes.

If necessary, adrenaline can be repeated at 5–15 minute intervals, while waiting for assistance.

**4. BE PREPARED TO COMMENCE AGE APPROPRIATE CPR\*** If needed.

**5. ADMINISTER OXYGEN** If available.

If there is respiratory distress, stridor, or wheeze, use high flow rates.

**6. RECORD VITAL SIGNS EVERY 5–10 MINUTES**

All observations and interventions need to be clearly documented in medical notes and should accompany the individual to hospital.

**7. ADMIT PATIENT TO HOSPITAL**

All cases of anaphylaxis should be admitted to hospital for observation.

Rebound anaphylaxis can occur 12–24 hours after the initial episode.

\*Note, a current Resuscitation certificate is required covering the skills outlined in Appendix 4.2 Immunisation Handbook.

