

# **National Immunisation Programme Operating Guidelines**

## **COVID-19**

## **Vaccines and General Operating Guidance**

Version 51.0

Last Updated February 2023

Citation: Ministry of Health. 2021. *Operating Guidelines for DHBs and Providers: COVID-19 Vaccine Immunisation Programme*. Wellington: Ministry of Health.

Published in September 2021 by the Ministry of Health  
PO Box 5013, Wellington 6140, New Zealand



This document is available at [health.govt.nz](https://health.govt.nz)



This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to: share i.e., copy, and redistribute the material in any medium or format; adapt i.e., remix, transform and build upon the material. You must give appropriate credit, provide a link to the licence, and indicate if changes were made.

## Section A: Ready to vaccinate - summary of changes

Version	Date	Section	Summary of Changes
51.0	28/02/23	Key contacts and Roles and responsibilities	Updated CIR contact details and hours Added hours for incident inbox Removed 'Group 1 Employers'
		2.1	Updated to add link to Te Whatu Ora website for current IPC advice.
		2.6	Reworded for clarity
		3.1-3.3	Updated to remove outdated information
		3.7	CIR drop in sessions are no longer available
		7.2	Removed reference to alert level settings, updated table 7.2 process step
		7.3/ table 7.3	Ordering site collateral has been updated with current guidance for how to register with BlueStar and what collateral is available to order/ download from Dropbox.
		Table 8.2	Syringe labels table updated with grey cap 30 mcg and 15/15 mcg. Vaccine packs have been updated for reference.
		Table 8.3	Vaccine shelf life table updated to reflect new grey cap vaccines.
		Table 9.1.5	Vaccine unit sizes and dimensions added for new grey cap vaccines, 10 mcg and 3 mcg Comirnaty vaccines have been updated.

## Section B: Pathway to COVID-19 vaccination - summary of changes

Version	Date	Section	Summary of Changes
51.0	28/02/23	17	Updated guidance for discontinued Comirnaty 30 mcg purple cap vaccine
		18-19	New sections with key safety points for Comirnaty 30 mcg grey cap and Comirnaty 15/15 mcg Original/Omicron BA.4/5 grey cap
		23	New section 'Preparation of doses' which summarises information previously found in each vaccine section.
		24.3	Removed 'my vaccine pass'

## Section C: Additional Programme guidance, variations and incidents - summary of changes

Version	Date	Section	Summary of Changes
51.0	28/02/23		No changes

## Appendices: summary of changes

Version	Date	Appendix	Summary of Changes
51.0	28/02/23	J	Updated table to remove outdated information

## Document approval

National Immunisation Programme	Date	Signature
Rachel McKay	28/02/23	Electronic

# Contents

National Immunisation Programme Operating Guidelines .....	1
<b>Introduction.....</b>	<b>9</b>
Purpose.....	9
Abbreviations .....	10
Key contacts.....	11
Roles and responsibilities .....	12
<b>Section A: Ready to vaccinate.....</b>	<b>13</b>
Section guidance .....	13
<b>1 Equity.....</b>	<b>14</b>
1.1 Equitable access .....	14
1.2 Te Tiriti and Māori .....	14
1.3 Māori and Pacific peoples .....	15
1.4 Disability and/or Impairments .....	15
<b>2 Site considerations.....</b>	<b>16</b>
2.1 Environmental considerations and safety controls at the vaccination site.....	16
2.2 Business continuity.....	16
2.3 Site access and traffic management.....	16
2.4 Site physical security.....	17
2.5 Planning for adverse events .....	18
2.6 Mobile vaccination set up .....	18
<b>3 Preparing the vaccination workforce.....</b>	<b>20</b>
3.1 Vaccinating the workforce.....	20
3.2 Clinical leadership.....	20
3.3 Preparation and planning phase.....	20
3.4 Quality and safety.....	20
3.5 Occupational health and safety requirements.....	20
3.6 Staff training and reference materials.....	21
3.7 Access to training, CIR classroom, and CIR.....	21
3.8 On site functions.....	22
3.9 Workforce modelling.....	22
3.10 Mobile and home vaccinator workforce .....	24
<b>4 Infection prevention and control (IPC) .....</b>	<b>25</b>

4.1	Key IPC principles for COVID-19 vaccine deployment .....	25
<b>5</b>	<b>COVID-19 Immunisation Register .....</b>	<b>27</b>
5.1	Logging in to CIR.....	27
5.2	Pre-loading immunisation event records in CIR.....	27
5.3	Where the consumer does not have an NHI number.....	28
5.4	Recording vaccine waste.....	28
5.5	CIR support .....	28
5.6	Recording in CIR.....	29
<b>6</b>	<b>Logistics .....</b>	<b>30</b>
6.1	Logistics.....	30
<b>7</b>	<b>Equipment ordering and demand planning.....</b>	<b>31</b>
7.1	Ordering IT equipment.....	31
7.2	Ordering personal protective equipment (PPE) .....	32
7.3	Ordering site collateral .....	32
7.4	Vaccine ordering/demand planning.....	34
7.5	Ordering Interwaste vial disposal bins.....	34
7.6	Ordering other NPHS Te Whatu Ora supplied consumables.....	35
7.7	CIR and inventory management .....	35
7.8	Operational reporting .....	36
<b>8</b>	<b>Vaccine storage and handling .....</b>	<b>36</b>
8.1	Vaccine security.....	36
8.2	Differentiation of vaccines.....	36
8.3	Cold chain storage .....	39
8.4	Movement of vaccine.....	42
8.5	Repacking vaccine at Health District facilities.....	42
8.6	Transportation of vaccine to other locations .....	42
8.7	Transportation of diluted or drawn-up vaccine .....	43
<b>9</b>	<b>Vaccine and consumables ordering and delivery .....</b>	<b>44</b>
9.1	Vaccine ordering .....	44
<b>10</b>	<b>Disposal of consumables, vaccine, and vaccine packaging.....</b>	<b>56</b>
10.1	Disposal of consumables .....	56
10.2	Disposal of damaged, empty, and expired vaccine vials .....	56
10.3	Disposal of vaccines drawn up but not administered and empty vaccine syringes.....	57
10.4	Vaccine packaging disposal.....	57

<b>11</b>	<b>Site readiness and closure.....</b>	<b>58</b>
11.1	Site setup form and site checklist.....	58
11.2	Completing a dry run .....	58
11.3	Facility/site closure form.....	58
11.4	Facility moving location.....	59
<b>12</b>	<b>Becoming a COVID-19 Vaccination site.....</b>	<b>60</b>
12.1	Onboarding.....	60
12.2	Additional resources.....	60
<b>Section B: Pathway to COVID-19 vaccination .....</b>		<b>61</b>
<b>13</b>	<b>Booking and scheduling.....</b>	<b>62</b>
13.1	Booking second doses.....	62
<b>14</b>	<b>Protecting security and privacy .....</b>	<b>64</b>
<b>15</b>	<b>COVID-19 vaccines operational phase .....</b>	<b>65</b>
<b>16</b>	<b>Obtaining informed consent.....</b>	<b>67</b>
16.1	Prescription .....	68
<b>17</b>	<b>Comirnaty purple cap COVID-19 vaccine (for ages 12 years and over) .....</b>	<b>69</b>
<b>18</b>	<b>Comirnaty 15/15mcg Original/ Omicron BA.4/5 grey cap COVID-19 vaccine (for ages 16 years and over) .....</b>	<b>70</b>
<b>19</b>	<b>Comirnaty 30mcg grey cap COVID-19 vaccine (for ages 12 years and over) .....</b>	<b>71</b>
<b>20</b>	<b>Comirnaty 10mcg orange cap COVID-19 vaccine (for ages 5 to 11 years) .....</b>	<b>72</b>
20.1	Site readiness .....	73
20.2	Vaccine safety and additional considerations for consumers aged 5 to 11 years.....	73
<b>21</b>	<b>Comirnaty 3mcg maroon cap COVID-19 vaccine (for ages 6 months to 4 years).....</b>	<b>74</b>
21.1	Vaccine safety and additional considerations for consumers aged 6 months to 4 years.....	74
<b>22</b>	<b>Nuvaxovid COVID-19 vaccine (for ages 12 years and over).....</b>	<b>75</b>
<b>23</b>	<b>Preparation of Doses.....</b>	<b>76</b>
<b>24</b>	<b>COVID-19 vaccine pathway to vaccination .....</b>	<b>79</b>
24.1	Sharing information on the vaccine.....	84
24.2	Observation following vaccination.....	84
24.3	Consumers' record of vaccination.....	85

<b>Section C: Additional Programme guidance, variations and incidents.....</b>	<b>86</b>
<b>25 Vaccination in high-risk or screened ‘positive’ consumers .....</b>	<b>87</b>
<b>26 Third primary dose for severely immunocompromised.....</b>	<b>90</b>
<b>27 Vaccination and Surveillance Testing .....</b>	<b>91</b>
<b>28 Vaccination in Hospital .....</b>	<b>92</b>
28.1 Introduction .....	92
28.2 Screening .....	92
<b>29 Mobile vaccination team.....</b>	<b>93</b>
29.1 Setting up mobile vaccination teams.....	93
29.2 Setting up in CIR .....	93
<b>30 Home vaccinations.....</b>	<b>94</b>
30.1 Transportation of vaccine for household vaccinations.....	94
30.2 Consumer Considerations .....	95
<b>31 COVID-19 Trial Vaccinations.....</b>	<b>95</b>
<b>32 National Immunisation Booking System .....</b>	<b>96</b>
32.1 Introduction .....	96
32.2 Booking system principles.....	96
32.3 Book my Vaccine system roles .....	97
32.4 Booking system processes and best practice .....	99
32.5 Setup and maintenance .....	99
<b>33 Incidents .....</b>	<b>103</b>
33.1 Incident management.....	103
33.2 Adverse events during observation period.....	103
33.3 Recording an anaphylaxis event .....	103
33.4 Adverse events after observation period.....	104
33.5 COVID-19 treatment injury claims .....	104
33.6 Recording vaccine errors .....	105
33.7 Early second doses.....	106
<b>34 Variations .....</b>	<b>107</b>
34.1 Missing or incorrect information in the CIR .....	107
34.2 Where the consumer has received vaccination overseas.....	107
<b>Appendices .....</b>	<b>108</b>

Section guidance .....	108
<b>Appendix A: Site checklist .....</b>	<b>109</b>
<b>Appendix B: New facility/site setup .....</b>	<b>115</b>
<b>Appendix D: Logistics and inventory management.....</b>	<b>122</b>
<b>Appendix E: NIP logistics overview/cheat sheets.....</b>	<b>123</b>
<b>Appendix F: Links to the National Immunisation Booking System.....</b>	<b>125</b>
<b>Appendix G: Vaccination site screening questions .....</b>	<b>126</b>
<b>Appendix H: Supported decision-making process.....</b>	<b>127</b>
<b>Appendix I: NIP Adverse Event Process .....</b>	<b>128</b>
Provider and Programme Lead Clinicians .....	129
Process Steps .....	130
<b>Appendix J: Risk mitigations for vaccination sites .....</b>	<b>134</b>
<b>Document version control .....</b>	<b>135</b>



# Introduction

These Operating Guidelines provide guidance on establishing and managing a COVID-19 vaccination site, including guidelines for the vaccination workforce and how to provide a clinically safe and quality vaccination service.

## Purpose

The Operating Guidelines are designed to assist Health Districts and providers to maintain public safety and to ensure consistent and equitable COVID-19 vaccination practices are established and maintained throughout Aotearoa New Zealand. The Operating Guidelines are to be read and interpreted in conjunction with the **Aotearoa New Zealand COVID-19 Vaccine Immunisation Service Standards (the Standards)**.

The Operating Guidelines are published on the **Ministry of Health's website** for Health Districts and providers. We expect regular iterations based on learnings from the delivery of the COVID-19 vaccine Programme. Please ensure the most updated version is used.

### Notes on guidance:

- The Operating Guidelines provide operational guidance for the COVID-19 vaccination Programme. Clinical guidance is available in the Immunisation Handbook, available at: <https://www.health.govt.nz/publication/Immunisation-Handbook-2020>.
- See in particular **Chapter 2 Processes for Safe Immunisation** and **Chapter 5 Coronavirus disease (COVID-19)**.

## Whakatauki

Me mahi tahi tātou mō te oranga o te katoa

*We should work together for the wellbeing of everyone*

# Abbreviations

Abbreviation	Full Name
<b>A&amp;I</b>	Adoption and Improvement
<b>AEFI</b>	Adverse Event Following Immunisation
<b>AIR</b>	Aotearoa Immunisation Register
<b>CARM</b>	Centre for Adverse Reactions Monitoring
<b>CICS</b>	COVID-19 Immunisation Consumer Support
<b>CIR</b>	COVID-19 Immunisation Register
<b>DNS</b>	Did not show
<b>IMAC</b>	Immunisation Advisory Centre
<b>IPC</b>	Infection prevention and control
<b>Ministry</b>	Ministry of Health
<b>NHI number</b>	National Health Index number
<b>NIBS</b>	National Immunisation Booking System (Book my Vaccine)
<b>NIP</b>	National Immunisation Programme (the Programme)
<b>NPHS</b>	National Public Health Service
<b>ULT</b>	Ultra-low temperature (-90°C to -60°C)

## Key contacts

Issue Type	When to Contact	Contact Details	Hours of Operation
<b>IT hardware or non-CIR software issues</b>	Logging technology hardware or software issues that aren't CIR-related	Contact your local IT ServiceDesk	Ensure after-hours support is available for sites operating outside of business hours
<b>CIR issues</b>	For help on using CIR Logging-in issues, password resets, or after hours help	Refer to the <i>Where to get help</i> poster* Email: <a href="mailto:help@C-19imms.min.health.nz">help@C-19imms.min.health.nz</a> Call: 0800 223 987	8am-5pm, Monday to Friday (from 9.30am on Wednesdays) 9am-2pm, Saturday and Sunday
<b>Vaccine or consumables supply issues</b>	To raise an issue with supplies	Refer to the <i>Where to get help</i> poster*	Email: 9am-5pm, weekdays
<b>Clinical vaccine queries</b>	To receive clinical advice on the vaccine or vaccination process	<b>0800 IMMUNE (466 863)</b> , option 1 (health professionals) and then option 2 (COVID-19 vaccinator support)	Available during site operating hours
<b>Order vaccination collateral</b>	To request additional pamphlets or other collateral	The Health District communications manager	
<b>Privacy Incident or Concern</b>	In the event of a known or suspected privacy breach	Refer to the <i>Where to get help</i> poster*	9am-5pm weekdays
<b>Adverse Event Following Immunisation (AEFI)</b>	Reporting an adverse reaction to the vaccine	<a href="mailto:report.vaccine.covid19.govt.nz">report.vaccine.covid19.govt.nz</a> Phone: (03) 479 7247 Email: <a href="mailto:carmnz@otago.ac.nz">carmnz@otago.ac.nz</a>	
<b>Interwaste vial disposal bin requests/collection</b>	To arrange first delivery of vial disposal bin and collection of full bins	Phone: 0800 102 131	8am-5pm, weekdays
<b>Programme Incidents</b>	See serious adverse event process <b>Appendix I</b>	<a href="mailto:nip.incidentnotification@health.govt.nz">nip.incidentnotification@health.govt.nz</a>	Inbox monitored 8.30am-4pm, weekdays

\*A **Where to get help** poster is available in the Ministry's drop box for vaccination sites. The poster includes the CIR helpdesk number and email address details, and NPHS Te Whatu Ora logistics team's contact number and email address.

# Roles and responsibilities

Activity	NPHS Te Whatu Ora	Health Districts & Providers	IMAC	CARM	Distribution Provider
<b>Purchasing</b>	<ul style="list-style-type: none"> <li>• Purchase vaccine</li> <li>• Purchase consumables including PPE</li> </ul>	N/A	N/A	N/A	N/A
<b>Distribution</b>	<ul style="list-style-type: none"> <li>• Arrange distribution of vaccine and consumables to vaccination sites/Health District facilities</li> </ul>	<ul style="list-style-type: none"> <li>• If needed, arrange secure distribution from Health District facility to vaccination site</li> </ul>	N/A	N/A	<ul style="list-style-type: none"> <li>• Thaw and repack vaccine into sub-batches as needed</li> <li>• Distribute vaccine &amp; consumables</li> </ul>
<b>Inventory Management</b>	<ul style="list-style-type: none"> <li>• Coordinate allocation schedule</li> <li>• Order vaccine &amp; consumables for Health Districts</li> </ul>	<ul style="list-style-type: none"> <li>• Plan vaccine demand to minimise wastage</li> <li>• Report stock on hand, stock movement &amp; exceptions</li> <li>• Ensure vaccine handling &amp; storage requirements are met</li> </ul>	N/A	N/A	<ul style="list-style-type: none"> <li>• Perform QA checks on receipt of vaccine from Pfizer</li> <li>• Ensure secure storage of vaccine prior to distribution</li> </ul>
<b>Workforce &amp; Training</b>	<ul style="list-style-type: none"> <li>• Provide guidance on workforce model and training requirements</li> <li>• Provide access to CIR for vaccinators &amp; admin staff</li> <li>• Provide CIR support/factsheets</li> </ul>	<ul style="list-style-type: none"> <li>• Hire and roster vaccinators and required site support staff</li> <li>• Provide info to NPHS Te Whatu Ora and IMAC for user on-boarding &amp; provision of training</li> <li>• Ensure staff are appropriately trained</li> </ul>	<ul style="list-style-type: none"> <li>• Provide vaccine preparation &amp; delivery training</li> <li>• Provide CIR training</li> </ul>	N/A	N/A
<b>Site Operations</b>	<ul style="list-style-type: none"> <li>• Provide guidance on preparing and running vaccination sites</li> <li>• Disseminate process improvements (e.g., via updated Operating Guidelines)</li> </ul>	<ul style="list-style-type: none"> <li>• Prepare &amp; run vaccination sites, incl. providing IT equipment and disposing waste</li> <li>• Work with Group 1 employers to schedule vaccinations of staff</li> <li>• Schedule appts for household contacts</li> <li>• Engage with Māori &amp; Pacific Island partners around vaccination of household contacts</li> </ul>	<ul style="list-style-type: none"> <li>• Provide clinical support to vaccinators as needed</li> </ul>	N/A	N/A
<b>Post-Event</b>	<ul style="list-style-type: none"> <li>• Monitoring adverse event data</li> </ul>	<ul style="list-style-type: none"> <li>• Dispose of expired, empty, or broken vaccine vials and used consumables</li> <li>• Pack down site as needed</li> </ul>	N/A	<ul style="list-style-type: none"> <li>• Receive and investigate adverse event reports</li> </ul>	N/A
<b>Comms &amp; Engagement</b>	<ul style="list-style-type: none"> <li>• Coordinate national vaccine engagement campaign</li> <li>• Provide key messages to Health Districts to share with Group 1 employers</li> <li>• Engage with household contacts</li> <li>• Provide collateral files to Health Districts/providers &amp; distribute site banners/cards</li> <li>• Manage adverse event comms</li> </ul>	<ul style="list-style-type: none"> <li>• Engage with Group 1 employers re: sites &amp; schedule</li> <li>• Print and circulate collateral to vaccination sites as required</li> <li>• Engage with household contacts</li> </ul>	N/A	N/A	<ul style="list-style-type: none"> <li>• Include 'Instructions for the Pfizer vaccine – Preparation and Administration' info sheet in vaccine shipments</li> </ul>
<b>Reporting</b>	<ul style="list-style-type: none"> <li>• Produce Programme and operational reporting</li> </ul>	<ul style="list-style-type: none"> <li>• Complete weekly stock on hand and stock movements reporting</li> <li>• Report exceptions to plan, as they occur</li> </ul>	<ul style="list-style-type: none"> <li>• Provide data on vaccinators trained to date</li> </ul>	<ul style="list-style-type: none"> <li>• Provide adverse event data to Medsafe</li> </ul>	<ul style="list-style-type: none"> <li>• Provide stock on hand and orders out reporting to NPHS Te Whatu Ora</li> </ul>

# Section A: Ready to vaccinate

## Section A: Ready to vaccinate - summary of changes

Version	Date	Section	Summary of Changes
51.0	28/02/23	Key contacts and Roles and responsibilities	Updated CIR contact details and hours Added hours for incident inbox Removed 'Group 1 Employers'
		2.1	Updated to add link to Te Whatu Ora website for current IPC advice.
		2.6	Reworded for clarity
		3.1-3.3	Updated to remove outdated information
		3.7	CIR drop in sessions are no longer available
		7.2	Removed reference to alert level settings, updated table 7.2 process step
		7.3/ table 7.3	Ordering site collateral has been updated with current guidance for how to register with BlueStar and what collateral is available to order/download from Dropbox.
		Table 8.2	Syringe labels table updated with grey cap 30 mcg and 15/15 mcg. Vaccine packs have been updated for reference.
		Table 8.3	Vaccine shelf life table updated to reflect new grey cap vaccines.
		Table 9.1.5	Vaccine unit sizes and dimensions added for new grey cap vaccines, 10 mcg and 3 mcg Comirnaty vaccines have been updated.

## Section guidance

This section should be read and interpreted in conjunction with **the Standards**.

This section provides operational guidance, including equity, site considerations, onboarding, vaccination workforce, IPC guidance, ordering, planning, vaccine handling and storage, logistics, and site closure; to ensure consistent, equitable and quality vaccination.

### Purpose

This section is designed to be applicable from the preparation of a vaccination site (from the selection and setting up of a suitable site), through to the closing of a site.

#### Appendices relevant to this section

- **Appendix A: Site checklist**
- **Appendix B: New facility/site setup**
- **Appendix C: Facility/site closure**
- **Appendix D: Logistics and Inventory Management**
- **Appendix E: NIP logistic overview/ cheat sheets**

# 1 Equity

Providers must ensure vaccination sites are accessible to all members of the community and there is equitable opportunity for Māori and Pacific people, other ethnic communities, and disabled people.

## 1.1 Equitable access

Reasonable steps must be taken to improve access and reduce potential inequalities. Steps to enable equitable access may include:

- Providing access to translation and interpretation services to support the consent and immunisation processes. For more information on interpreter services see <https://www.healthnavigator.org.nz/languages/i/interpreter-services/>
- Ensuring key written material and any signage is in easy-to-read formats.
- Providing supporting literature available in a range of languages and resources/support for those who have low health literacy. This may include access to New Zealand Sign Language (NZSL) if needed.
- Considering how the service delivery model caters for the support people consumers may bring to the vaccination event (such as friends, whānau, carers).
- Encouraging site staff to greet consumers in Te Reo or the language the consumer uses where possible.

## 1.2 Te Tiriti and Māori

Actively incorporate Te Tiriti o Waitangi considerations, including:

- ensuring Māori are not disadvantaged
- mitigating the impact to Māori as a result of COVID-19
- establishing and maintaining effective partnerships with Māori stakeholders including iwi, hapū and whānau
- seeking Māori-specific advice from the outset
- resourcing and investing where it is required the most
- starting and ending the day with a karakia.

## 1.3 Māori and Pacific peoples

- Ensure as far as reasonably practicable, the site workforce reflects the demographic make-up of the likely consumer group or local area.
- Consider which site locations can best meet the community's needs in terms of both ease of access and comfort or familiarity with the location (such as marae, churches).
- Where drive-in sites are planned, ensure consumers can attend the site if they do not have a car or have access to a non-drive-in site.
- Build early and regular engagement with Māori and Pacific partners into the service delivery model to ensure design to the community's needs.

## 1.4 Disability and/or Impairments

Ensure access for disabled consumers and others, including venue accessibility and accessible information. For more information on venue accessibility, see the **Ministry's website**. Equity steps and processes to follow include:

- Designing site support processes to support consumers with visual impairments or are hard of hearing. For example, providing a card to ask consumers advise site staff if they have a hearing impairment to ensure their needs can be met during the vaccination or any follow up interactions.
- For Deaf or hard of hearing consumers, there may be a need to arrange a New Zealand Sign Language (NZSL) Interpreter. Information on working with NZSL Interpreters can be found at <https://www.odi.govt.nz/nzsl/tools-and-resources/>
- Ensuring staff are educated in disability equity issues and know how to employ a rights-based approach. A 30-minute Disability Equity eLearn is available through the **Ministry's LearnOnline website**.
- Enabling consumers to access appropriate support and accommodations they may need for a successful vaccination, for example, are there any measures as a site or team that can be implemented to support mobility constraints, or accommodate individuals, families and whānau if a consumer has an anxiety or phobia, or may need a quiet and low stimulation environment?
- Supported decision-making is an important process for consumers needing support to make decisions. This may be due to a consumer's communication needs, learning disability, acquired brain injury, neurodiverse needs, mental health issues or other cognitive or physical condition.
- Supported decision-making is a way for consumers to make their own decisions based on their will and preferences, so they have control of their life, ensuring the consumer needing support is at the centre of decision making that concern them. Training on supported decision making is available on **IMAC's website**.

## 2 Site considerations

### 2.1 Environmental considerations and safety controls at the vaccination site

Assess the layout of the building or area identified for vaccination delivery to ensure features are in place supporting appropriate IPC implementation to meet current required standards for the site location. This is to ensure protection for consumers as well as site staff.

For current advice, refer to [COVID-19: Infection prevention and control recommendations for health and disability care workers – Te Whatu Ora - Health New Zealand](#).

### 2.2 Business continuity

A business continuity plan is required for each site to guide recovery from events that may interrupt service delivery such as a power failure.

Hard copies of the following forms and documents should be available on site in the event of the CIR being unavailable:

- **Consent form** (required consumer data fields that will need to be added to CIR are included on the back of the form)
- **COVID-19 Vaccine Adverse Event Report form.** This form is used to submit adverse event information to the Centre for Adverse Reactions Monitoring (CARM). If CIR is unavailable this form may be used to capture relevant information; noting on-site adverse events must in any event be reported in CIR as soon as practicable (in addition to submitting the form to CARM).
- **Reviewing early COVID-19 AEFIs** (found on the **IMAC website**)
- **Reviewing late onset AEFIs** (found on the **IMAC website**)

See the **Ordering site collateral** section below for obtaining these forms.

**Note:** Any hard copy forms must be entered into CIR as soon as practicable and in any event by close of business on the **following day**. Ensure any printed copies of information are locked away when not in use.

### 2.3 Site access and traffic management

Waka Kotahi NZ Transport Agency has provided the following advice to support site location and traffic management planning.



In addition to the considerations below, the **Waka Kotahi Journey Planner** is useful for assessing how people will safely access your sites. Similarly, regional council websites also contain valuable information about local public transport provision.

## Access considerations

When choosing your location, consider how easily people might be able to access the site. For example, consider the following:

- How easily people with mobility issues can access your site
- Is a public transport stop within 500m of your site?
- Are there multiple routes and/or multiple modes of public transport within 500m?
- Does the site provide cycling or walking access?
- Is adequate parking available for people using a private vehicle?
- Are there opportunities to locate the site in place that will reduce the number of additional trips people need to make?
- Is any additional signage required to direct people to the location of the centre?
- How would consumers living in areas not serviced by public transport reach your site?

## Traffic management considerations

Consider how the numbers of people receiving vaccines increases will impact the traffic network. For example, consider:

- How will the increase in road users impact vehicle congestion?
- How many different routes can consumers use to access the site?
- The impact to current levels of congestion at different times of the day.
- Is the site close to major arterial roads or state highways, which may give greater access?
- Does your site location provide easy access to public transport to mitigate impacts on road congestion?
- Are there any planned roadworks, road closures, or events that may impact access?
- Will any potential queues to your facility affect access to key services such as emergency services, health centres or schools?
- Could you provide multiple small sites instead of a few major locations servicing large numbers of people to better disperse demand across the transport system?
- Can your booking system be used to manage demand on the facility and consider peak traffic times?

## 2.4 Site physical security

To ensure the safety of consumers and staff, all vaccination sites should consider a security presence to control access and to be available to support in the event of attempted unauthorised access, or protest action.

Staff should have access to safe accessible parking or public transport as required.

### Site security assessment

Each vaccination site must provide for:

- Staff safety

- Consumer safety
- Visitor safety
- Vaccine security including storage facilities and in-transit
- Information security – particularly paper-based information such as spreadsheets
- Contingency plans addressing a disturbance/potential protest event.

A documented risk assessment should be conducted for every individual vaccination site. This should include, but is not limited to, the following considerations:

- How will staff travel to the vaccination location?
- Will secure parking be provided for vaccinators and administrators?
- How is site access controlled?
- How is the vaccine transported to and from the vaccination site?
- How is the vaccine securely stored at the vaccination site?
- How are consumables, including items such as needles, securely stored at the vaccination location?
- How is hard copy information (if any) securely stored at the vaccination site?
- How staff respond to disruptions (such as protest action).

## 2.5 Planning for adverse events

Consumers who have a history of allergy or hypersensitivity, following administration of vaccines or injectable medicines, will require additional monitoring at the time of receiving their first vaccine dose. Similarly, consumers who experienced an adverse event after receiving their first dose of the vaccine may require clinical monitoring at the time of the second dose.

NPHS Te Whatu Ora expects vaccination sites to have appropriate protocols, equipment, settings, and workforce in place to support those who may require enhanced care following vaccination. Consider arranging any enhanced or additional consumer care requirements at the time of booking, or prior to these consumers attending a vaccination site.

It is recommended simulation scenarios are used to prepare staff to respond to adverse events.

## 2.6 Mobile vaccination set up

Mobile vaccination teams may be established to reach vulnerable families or small communities to address equity needs for the community being vaccinated.

When setting up a mobile vaccination team, it is important an appropriate operating model is in place and includes the following:

- **Equipment and connectivity:** Ensure mobile vaccination teams have the required equipment, both medical equipment and technology, to enable the use of CIR onsite. Check the connectivity at the site before attending.
- **CIR recording:** Ensure the mobile team know the name of their facility and team (site) to select in CIR.
- **Planning:** Establish a location plan for the mobile team with the appropriate logistics in place. Ensure a record is kept of where and when the mobile team has been

vaccinating and notification to local services as required eg emergency services, local iwi etc.

- **Vaccine storage and transport:** All appropriate and standard cold chain requirements must be met when transporting and storing vaccine. See guidance on transporting and storing vaccine in the **Vaccine storage and handling** section below for more information.
- **Business continuity:** Ensure a business continuity plan is in place for the team to manage unexpected events and appropriately record vaccination events, such as having a stock of printed event forms on hand if access to CIR is unavailable and managing unexpected events such as protest action.
- **Site readiness:** Refer to the **Site readiness and closure** section below for completing a dry run with your mobile team before commencing vaccinations.

# 3 Preparing the vaccination workforce

## 3.1 Vaccinating the workforce

Before commencing vaccinations, the Programme recommends all staff are provided with an opportunity to ensure they are up to date with all vaccinations including any eligible COVID-19 doses.

## 3.2 Clinical leadership

Every vaccinating site or service should have a named lead clinician each shift. This lead should be an appropriately experienced clinician who is able to lead the vaccination team, manage and investigate adverse events and incidents, and provide onsite clinical advice.

## 3.3 Preparation and planning phase

- Appoint an appropriate team member as the IPC lead for the service.
- Identify an adequate number of vaccinators and administrators to ensure sufficient staff and time is available to support correct implementation of IPC practices required to cover any staff absences and provide for consumers requiring any extra support.

## 3.4 Quality and safety

There is an expectation that each District has quality and safety oversight of the vaccination Programme rollout through their existing quality and safety and/or clinical governance mechanisms. For clarity, this includes adverse events, complaints, and incident management.

**Note:** In this context, 'adverse event' does not refer to an adverse reaction following immunisation.

## 3.5 Occupational health and safety requirements

Appropriate occupational health and safety policies and procedures are required for each site. This will include an accessible needlestick injury protocol which staff are familiar with.

## 3.6 Staff training and reference materials

Training will be provided to CIR users and vaccinators through a combination of eLearning Modules and quick step guides. The quick step guides will be available within the eLearning system, as well as within the knowledge tab of the CIR for continued availability and reference.

The eLearning modules and quick step guides include:

- Working with the COVID-19 Immunisation Register (eLearning)
- COVID-19 vaccinator education course (eLearning)
- COVID-19 vaccination for prescriber health professionals (eLearning)
- CIR quick step guides: reception, vaccination, recovery, quick adverse event, adverse event
- Inventory management (eLearning)

In addition to these training materials, staff have access to a range of reference materials. Please refer to the IMAC website for vaccinator training materials. These include:

- IMAC written resources: <https://covid.immune.org.nz/faq-resources/written-resources>. This includes COVID-19 vaccinator guidelines and instructions for preparing doses.
- IMAC video resources: <https://covid.immune.org.nz/faq-resources/video-resources>
- IMAC FAQs: available on the IMAC website at: <https://covid.immune.org.nz/faq>
- ***The Immunisation Handbook***: provides clinical guidance for administering vaccines. IMAC has also prepared a COVID-specific chapter in the Handbook. This information is updated regularly. See <https://www.health.govt.nz/publication/immunisation-handbook-2020>

See the **Ordering site collateral** section below for details regarding collateral to be given to consumers.

## 3.7 Access to training, CIR classroom, and CIR

Staff are required to complete the IMAC training by registering at [lms.immune.org.nz](https://lms.immune.org.nz). Users will complete CIR and/or Pfizer eLearning modules.

To support their training, CIR users will be granted access to CIR classroom to practice using the system. To gain access to CIR classroom, the Health District or provider workforce lead must send a list of all staff requiring CIR Classroom access to NPHS Te Whatu Ora.

Once staff have completed the required training, the Health District or provider workforce lead must confirm to the Programme that the staff member is 'approved'; The Programme will then provide access to the live CIR environment.

**Note:** An organisation email address must be supplied for any CIR user to obtain access to the live CIR environment.

## 3.8 On site functions

NPHS Te Whatu Ora has identified the following functions for the onsite team. Note that someone with a clinical role (such as a vaccinator) may perform non-clinical functions, particularly in smaller sites.

The list below outlines the functions required to assist workforce planning. It is not intended to be a prescriptive list of all functions and expectations of different roles.

### Clinical functions

- Preparing the vaccination dose
- Obtaining consent to receive the vaccination
- Asking health questions prior to administering the vaccine
- Vaccinating the consumer
- Monitoring consumers in an observation area for any adverse events
- Attending to adverse events and recording them

Staff performing clinical functions must be appropriately trained by **the Immunisation Advisory Centre (IMAC)**.

### Non-clinical functions

- Greeting consumers and answering questions
- Identifying any accommodations and additional support consumers may require, such as mobility support, low sensory/quiet spaces, interpreters (including New Zealand Sign Language interpreters)
- Confirming consumer identity
- Entering consumer information into CIR
- Ensuring the up-to-date consumer collaterals are in stock including consent form and vaccine information fact sheets
- Directing the consumer to the Privacy Statement
- Recording the vaccine details in CIR
- Advising the consumer when they can depart the observation area
- Completing or arranging daily cleaning of the site
- Arranging collection of medical waste
- Decommissioning the site when it is no longer needed
- Providing reporting back to NPHS Te Whatu Ora or Health District or provider leads as needed.

## 3.9 Workforce modelling

The size of the vaccination site and volume of vaccinations expected to be delivered on site will determine the size of the workforce required. The following tables outline staffing models for consideration as the vaccination workforce is planned.

**Note:** The framework below is only a suggestion and site workforce requirements will depend on matters such as expected site volumes, the service delivery model adopted and the likely needs of the consumers (for example, low health literacy or low English skills), more support throughout the process may be required which may in turn affect timing and resourcing.

Refer to **Appendix 4** in the *Immunisation Handbook* for further guidance on criteria for authorised vaccinators and minimum staff and equipment requirements for the provision of vaccination services.

**Table 3.1 – activities and associated staffing**

Waiting room	Immunisation event	After the event
<p><b>Activity</b></p> <ul style="list-style-type: none"> <li>Consumer checked in to the site as well as the CIR, any additional support required by consumer is arranged.</li> </ul>	<ul style="list-style-type: none"> <li>Consumer and vaccinator will have a clinical conversation about the vaccination and consumer will provide consent</li> <li>Immunisation occurs</li> <li>Administrator will enter details into CIR as the vaccinator performs the vaccination</li> </ul>	<ul style="list-style-type: none"> <li>Consumers must remain onsite for 15 mins after the event for monitoring.</li> </ul>
<p><b>Staffing</b></p> <ul style="list-style-type: none"> <li>1 x Administrator</li> </ul>	<p><b>Staffing</b></p> <ul style="list-style-type: none"> <li>1 x Administrator</li> <li>1 x Vaccinator</li> </ul>	<ul style="list-style-type: none"> <li>1 x Registered health professional <b>minimum specifications in Appendix 4.2</b> of the <i>Immunisation Handbook</i>.</li> <li>1 x support person with CPR training</li> </ul>

Based on the activities and staffing numbers above, NPHS Te Whatu Ora recommends the following site staffing numbers:

**Table 3.2 – site staffing number recommendations**

If 20 vaccinations/day	If 120 vaccinations/day	If 360 vaccinations/day
<p><b>Staffing</b></p> <ul style="list-style-type: none"> <li><b>2 x vaccinators working at the site who will undertake all roles</b></li> </ul>	<p><b>Staffing</b></p> <ul style="list-style-type: none"> <li>1 x Admin in waiting room</li> <li>3 x Vaccinators</li> <li>3 x Admin support</li> <li>1 x Vaccinator drawing up</li> <li>1 Registered Health Professional and 1 x Support person monitoring during observative period</li> </ul>	<p><b>Staffing</b></p> <ul style="list-style-type: none"> <li>1 x Admin in waiting room</li> <li>9 x Vaccinators</li> <li>9 x Admin support</li> <li>3 x Vaccinators drawing up</li> <li>2 x Registered Health Professionals and 1 x Support person monitoring during observative period</li> </ul>

**Note 1:** If Vaccinating Health Workers are being used, there must be one (1) dedicated vaccination clinical supervisor for every six (6) Vaccinating Health Workers.

**Note 2:** Dedicated vaccination clinical supervisors are not simultaneously responsible for any other roles or processes that prevent them from being immediately available while supervising Vaccinating Health Workers.

**Note 3:** Health Districts and providers will need to be prepared to adjust their site staffing requirements as administering the COVID-19 vaccine will likely vary from these assumptions as delivery progresses and lessons learned

## 3.10 Mobile and home vaccinator workforce

For fixed sites, providers should consider the number of vaccinators and administrators that are needed for home or mobile vaccinations to ensure safety of both consumers and staff. Staff delivering home vaccination will need to meet the standards as set out in the **COVID-19 Vaccine and Immunisation Programme Service Standards** and have completed the required training.



# 4 Infection prevention and control (IPC)

For the latest Ministry guidelines on IPC please see the following [link](#). These principles and recommendations have been derived from the World Health Organization (WHO) guidance.<sup>1</sup>

This guidance is intended for policy makers, immunisation Programmes and IPC Lead for vaccination delivery venues.

## 4.1 Key IPC principles for COVID-19 vaccine deployment

Standard precautions to be applied during any vaccination activity are also valid for COVID-19 vaccine delivery, considering the population to be vaccinated consists of individuals **not** presenting signs and symptoms of infection.

National guidance and protocols for IPC measures should be consulted and adhered to when developing site operational guidelines.

### Local IPC guidance

Include the following details, when developing your local IPC guidance and standard operating procedures for COVID-19 vaccination:

- Screening policies for COVID-19 signs and symptoms for staff and consumers arriving for vaccination along with clear exclusion criteria.
- Key IPC measures to be taken by anyone in the vaccination area or clinic.
- Key IPC measures for safely administering COVID-19 vaccines.
- Cleaning and disinfection of the environment.
- Appropriate waste management, taking into consideration the increase of waste associated with COVID-19 vaccination activities. Where possible, include environmentally sound approaches to manage both general and medical waste at point of use, segregation, disposal, and collection.
- Visual reminders emphasising hand hygiene, safe injection practices, respiratory hygiene, and other IPC measures.
- Training materials for relevant staff.
- Communication material to inform and educate consumers.

---

<sup>1</sup> Aide-Memoire Infection prevention and control (IPC) principles and procedures for COVID-19 vaccination activities, 15 January 2021. <https://apps.who.int/iris/handle/10665/338715>

## IPC supplies

Ensure there is a continuous and sufficient supply of the following as required to conform to current guidelines:

- PPE, including eye protection and long-sleeve fluid resistant gowns and gloves for the vaccination team's protection in the event of dealing with a vaccine adverse event or other incidents such as support to an unwell consumer or clean-up of body fluids.
- Other IPC supplies including alcohol-based hand sanitisers, thermo-scans for temperature screening, tissues, waste bins and bin liners, sharps disposal bins, cleaning and disinfection products, visual reminders, and signage and physical barriers to aid spatial separation.

Identify a suitable secure area for storage of supplies.

# 5 COVID-19 Immunisation Register

The COVID-19 Immunisation Register (CIR) is a centralised, browser-based system used to record all vaccination details. CIR uses email address, phone number and six identifiers to match consumer records with NHI records.

Once a site has joined the National Immunisation Programme (NIP), request access to CIR for vaccinators and administrators, following the process outlined below.

For any questions or support on new user onboarding, the regional account manager should be contacted.

## 5.1 Logging in to CIR

Access request is made to the CIR location and inventory portal by contacting NPHS Te Whatu Ora, email [help@C-19imms.min.health.nz](mailto:help@C-19imms.min.health.nz) or call **0800 223 987**.

- To access to the live CIR Location and Inventory portal follow the link <https://ncts.force.com/cir/s/>
- Email [help@C-19imms.min.health.nz](mailto:help@C-19imms.min.health.nz) or call **0800 223 987**, for assistance with forgotten passwords or logging in problems.

Further details regarding how to log into the CIR can be found in the quick guides, videos, and detailed training guide at <https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Inventory-Orders-Portal-Quick-Step-Guide>.

## 5.2 Pre-loading immunisation event records in CIR

CIR is linked to consumers' NHI numbers, meaning any consumer with an NHI will automatically be available in CIR (they will have a CIR profile). Where consumers are in the border worker testing register (BWTR), NPHS Te Whatu Ora will extract that information to create immunisation event records (or cases) and add these to the consumer's CIR profile.

If consumers aren't in the BWTR, vaccinators or site administrators can add the immunisation event record/case to the consumer's profile on site at the time of vaccination.

## 5.3 Where the consumer does not have an NHI number

Where a consumer does not have an NHI in CIR, create a new NHI number for that consumer. If you do not have the ability to create an NHI number in Health UI, contact the Ministry contact centre on 0800 855 066 to request an NHI number be set up.

When contacting the centre:

- Provide the payee number for the Health District or hospital
- Identify the COVID-19 vaccination clinic
- Provide the name of the consumer
- Once the NHI is created, make sure it is linked to CIR using the NHI retrieval function. Retrieving the NHI will create a person profile in CIR which can then be used to create immunisation case records as normal.

**Note:** It is not mandatory to collect information on the consumer's residency status when setting up new NHI numbers. Experience has demonstrated that collecting residency information can be a barrier for consumers both in their uptake and receipt of healthcare services.

## 5.4 Recording vaccine waste

It is important for vaccine sites to record vaccine waste in the CIR Logistics Portal, but only to the unopened vial level (the recording of vaccine wasted at the opened vial level is yet to be determined). This is so that vaccine vial waste can be tracked at a local, regional, or national level.

Further details regarding how to log into the CIR can be found in the quick guides, videos, and detailed training guide at <https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Inventory-Orders-Portal-Quick-Step-Guide>.

See the Standard operating procedure (SOP) for inventory management on this [SOP for inventory management CIR link](#)

## 5.5 CIR support

If the site team requires CIR support, they should contact their super user in the first instance before contacting the CIR ServiceDesk.

CIR eLearning modules and quick step guides are available to all staff (see the [Staff training and reference materials](#) section above).

## 5.6 Recording in CIR

### CIR reports

The CIR portal provides a centralised place for operational reporting, including demand forecast, inventory management (including stock on hand), and orders approved for sites.

These operational reports can be generated for providers by the NPHS Te Whatu Ora NIP logistics customer services team and will be made available to providers in the future.

### Available hard copies

Hard copies of the following forms should be available on site, in the event of CIR being unavailable:

- **Consent forms**  
the required consumer data fields that need to be added to CIR are included on the back of the form
- **COVID-19 Vaccine Adverse Event Report**  
this is the form used to submit adverse event information to CARM. In the event of CIR being unavailable, this form can be used to capture relevant information, noting that on-site adverse events must be reported in CIR as soon as practicable (as distinct from submitting the form to CARM).

See the **Ordering site collateral** section below regarding obtaining these forms.

# 6 Logistics

## 6.1 Logistics

NPHS Te Whatu Ora will maintain the COVID-19 Immunisation Register (CIR) logistics module to support ongoing monitoring of inventory and demand. **Appendix D** shows the current process for distributing the vaccine to vaccination sites. **Appendix E** provides NIP logistics overview/ cheat sheets.

### Logistics support

NPHS Te Whatu Ora provides two levels of customer support.

- Level one is NPHS Te Whatu Ora's IT helpdesk. The helpdesk deals with log-in and access issues and can be contacted by emails: **help@C-19imms.min.health.nz** or by phone on **0800 223 987**.
- Level two is the NIP logistics customer services team. This team can assist with support for order placing and approval, inventory management, and use of the CIR inventory portal. Once the vaccination site has been onboarded, contact details for this team will be provided.

### Quality Assurance Approval Step of Orders

Supplier orders made by Inventory users at a Health District level will be sent to their Quality Assurance (QA) user to be reviewed and approved before being sent to Te Whatu Ora for approval. The QA user can add and remove products from the order as well as edit the quantity of these products in the order. The QA user can also reject the order or accept the order. Accepting the order will send it through to NPHS Te Whatu Ora Logistics team for approval. Each Health District and Provider using the inventory portal will need to have dedicated QA users to review these orders. If a supplier order is created by a QA user, it will go straight to NPHS Te Whatu Ora logistics team for approval.






Further detail about how to log into the CIR can be found in the quick guides, videos, detailed training guide on <https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Inventory-Orders-Portal-Quick-Step-Guide>

# 7 Equipment ordering and demand planning

## 7.1 Ordering IT equipment

Provide the IT requirements, outlined in table 7.1 below, at vaccination sites to ensure staff can access the COVID-19 Immunisation Register. Before starting vaccinations, ensure all IT equipment has been tested, and all staff have received the necessary training to use the devices and CIR. Advise each site team where they can access additional IT support (for non-CIR issues such as hardware issues), including after-hours support if your vaccination site is operating outside standard business hours.

**Table 7.1 – IT requirements**

Requirement	Details
 <b>Network</b>	<ul style="list-style-type: none"> <li>A secure network (Wi-Fi, hard wired, or 4G) with connectivity to the device running CIR, and to the user's mobile phone or computer.</li> <li>Site Wi-Fi specifications: Coverage ranging to reception, vaccination and waiting areas Highly available network (such as fibre and 4G backup)</li> </ul>
 <b>Internet Browser</b>	<ul style="list-style-type: none"> <li>Chrome is the recommended internet browser. Other browsers support CIR, but Internet Explorer is not supported (use Microsoft Edge if needed). For further information see: <a href="https://help.salesforce.com/articleView?id=sf.getstart_browsers_sfx.htm&amp;type=5">https://help.salesforce.com/articleView?id=sf.getstart_browsers_sfx.htm&amp;type=5</a></li> </ul>
 <b>Computer or Tablet Device</b>	<ul style="list-style-type: none"> <li>Any laptop from the last five years should be compatible with CIR providing it has the appropriate browser access.</li> <li>For further information see: <a href="https://help.salesforce.com/articleView?id=sf.getstart_browser_recommendations.htm&amp;type=5">https://help.salesforce.com/articleView?id=sf.getstart_browser_recommendations.htm&amp;type=5</a></li> </ul>
 <b>Mobile Phone</b>	<ul style="list-style-type: none"> <li>CIR users require an iOS or Android mobile phone to download the Salesforce Authenticator application. This can be downloaded from the App Store on iOS and the Play Store on Android.</li> <li>You can scan the QR code on the right to locate the Salesforce Authenticator app in the relevant App Store.</li> </ul> 

## 7.2 Ordering personal protective equipment (PPE)

Table 7.2 – information required when ordering PPE

Details	Process
<b>PPE provided will be based on the current COVID-19 COVID-19 IPC guidelines (see section 4 above)</b>	<ul style="list-style-type: none"><li>• Order via the existing PPE portal via HealthCare Logistics or Onelink</li><li>• If you are new or currently do not hold contingency stock, please contact <b>COVID.healthsupplychain@health.govt.nz</b> to discuss your requirements</li></ul>

## 7.3 Ordering site collateral

The Programme has prepared collateral to support the vaccination programme. Files will be shared via the NIP Dropbox or via a Ministry weblink. You can download these files and print locally as required. Please ensure you are using the latest version of collateral.

Hardcopies of some resources can be ordered via the Blue Star portal.

To register with the Blue Star portal, email [moh.support@bluestargroup.co.nz](mailto:moh.support@bluestargroup.co.nz) and request 'Vaccine resources access'; include the following details:

- Organisation Name
- Name of person
- Email
- Phone
- Physical address

Once the person has been set up as a user. They will receive an email from Bluestar with log on details. Access can take a few working days.

IMAC has also prepared a consent video which can be displayed in site reception areas if desired. This video is available on the **IMAC website**.

**Note:** An interpreter may be arranged to be available on site to assist consumers who speak languages other than English, including New Zealand Sign Language. See the **Equitable access** section above for more information about interpreters.

Some collateral items have been translated and are available on Dropbox and/or Blue Star.



**Table 7.3 – site collateral ordering and purpose**

<b>Collateral</b>	<b>Purpose</b>	<b>How to Order</b>
<p><b>COVID-19 informed consent suite, which includes:</b></p> <ul style="list-style-type: none"> <li>• <b>HP8590 What you need to know about the COVID-19 vaccination</b></li> <li>• <b>HP7565 COVID-19 vaccination consent form</b></li> <li>• <b>HP8591 After the COVID-19 vaccination</b></li> <li>• <b>HP7568 Privacy statement</b></li> </ul>	To share with consumers on site or before attending the vaccination site	<p>Download directly from the NIP Dropbox.</p> <p>Some items are available in hard copy via Blue Star portal.</p>
<b>COVID-19 vaccine FAQs</b>	To provide answers to FAQs	Available on the <b>Ministry’s website</b>
<b>Booster collateral</b>	To provide information on COVID-19 boosters	<p>Download directly from the NIP Dropbox.</p> <p>Some items are available in hard copy via Blue Star portal.</p>
<b>Consent form (which includes fields to capture required consumer data)</b>	For use if CIR is unavailable	Download directly from the NIP Dropbox.
<b>COVID-19 Vaccine Adverse Event Report</b>	To provide information, and to enable accurate record keeping	Available on the Centre for Adverse Reactions Monitoring (CARM) website: <a href="https://report.vaccine.covid19.govt.nz">https://report.vaccine.covid19.govt.nz</a>
<b>Vaccine Error Reporting Form</b>	To enable accurate record keeping	Contact the Health District communications manager
<b>Vaccination station material</b>	Promotional material to support vaccination site set up	Order via Blue Star
<b>Instructions for the Preparation and Administration of vaccines</b>	For vaccinators and staff on site	Included in vaccine shipments and are available on the <b>IMAC website</b> .
<b>‘Where to get help’ poster</b>	To provide information simply and quickly	<ul style="list-style-type: none"> <li>• Contact the Health District communications manager</li> <li>• Also available via the CIR homepage</li> </ul>

## 7.4 Vaccine ordering/demand planning

Table 7.4 – site and facility set up for vaccine delivery

Information required	Details	Process
<b>Site and facility set up information</b>	<ul style="list-style-type: none"> <li>Site and facility information must be provided to NPHS Te Whatu Ora five (5) days in advance of any initial deliveries.</li> </ul>	<ul style="list-style-type: none"> <li>Use the <b>New facility site set up form (found in Appendix B)</b> to submit site or facility details</li> <li>Return the completed form via email to <b>help@c-19imms.min.health.nz</b></li> </ul>

Table 7.5 – demand planning

Information required	Details	Process
<b>Demand plan – appropriate to cater for the upcoming four weeks</b>	<ul style="list-style-type: none"> <li>The plan should represent the expected number of vials to be consumed each day, in each location, for the upcoming four-week period.</li> <li>The plan should be maintained at the facility level on the vial's product.</li> <li>21 days of demand forecast must be loaded for a location to place an order.</li> </ul>	<ul style="list-style-type: none"> <li>Upload and maintain the plan in the CIR Inventory Portal using the demand upload functionality.</li> <li>Please update forward forecasts on a weekly basis.</li> </ul>

## 7.5 Ordering Interwaste vial disposal bins

As part of site preparations, Interwaste must be contacted to arrange the delivery of an Interwaste vial disposal bin (see the **Disposal of consumables, vaccine, and vaccine packaging** section below).

Contact Interwaste on 0800 102 131 (business hours) as soon as the site is approved. Provide at least five business days' notice before the container is required to arrive. Interwaste will collect the relevant details such as the site manager's name and contact details, the delivery date for the first container, and the site delivery address information.

## 7.6 Ordering other NPHS Te Whatu Ora supplied consumables

Table 7.6 – Other consumables

Information required	Details	Process
<b>Order for other consumables (such as sharps bins, bio bags for waste disposal, or 21G 38mm needles)</b>	<ul style="list-style-type: none"> <li>This stock will be shipped through a standard courier network, expect delivery between two and four days from the time of order.</li> </ul>	<ul style="list-style-type: none"> <li>Order consumables via the CIR Portal.</li> </ul>
<b>Order for other individual items (such as boxes of plasters).</b>	<ul style="list-style-type: none"> <li>This stock will be shipped through a standard courier network, expect delivery between two and four days from the time of order.</li> </ul>	<ul style="list-style-type: none"> <li>Order consumables via the CIR Portal.</li> </ul>

## 7.7 CIR and inventory management

The COVID-19 Immunisation Register (CIR) provides a centralised place for vaccine and consumables orders, managing stock on hand (SOH), arranging transfers, and recording consumption and wastage of unopened vaccine vials.

The CIR inventory module is where movement (transactions) and use of stock is managed and recorded. (The term inventory is used to describe how much product or stock (in this case vaccine and consumables) is at a location at any point in time.) These records provide effective stock management at each location, ensuring optimum use – and minimum wastage – of vaccines and consumables.

NPHS Te Whatu Ora logistics team will continue to monitor demand and allocation using data from CIR along with information provided by Health Districts or providers. Health District and provider logistics leads must supply daily reporting (as required) on:

- Stock on hand (daily stock takes)
- Stock movements, including ordering, transfers, wastage, consumption, and stock adjustments
- Stock consumption
- Stock waste
- Quarantine of and repacking of stock.

NPHS Te Whatu Ora logistics team will liaise with logistics leads to collect this information through an agreed mechanism.

Health Districts or providers may wish to collate daily reporting back from sites on inventory and/or operations to aid the supply of information back to the NPHS Te Whatu Ora.

Please contact your regional liaison if you have feedback on the immunisation process or recommendations for operational improvements.

## 7.8 Operational reporting

Health Districts or providers need to report significant events on sites such as a significant adverse reaction, or a protest to the Ministry on a daily basis.

# 8 Vaccine storage and handling

## 8.1 Vaccine security

To ensure the security of the vaccine, the following minimum standards must be met:

- Vaccines must be stored in a work area that has the constant presence of an authorised person (such as an administrator, or security guard or vaccinator) during the hours of operation.
- If the vaccine is to be stored overnight at the vaccination site, the building should be in a controlled-access environment (such as Maritime Port).
- If the building is not in a controlled-access environment (such as a community hall), the building should be able to be secured and have a monitored alarm.
- In the event of the vaccines being stored at a vaccination site without controlled access and not a building (such as a tent), an overnight onsite security guard must be present.

## 8.2 Differentiation of vaccines

Syringe labels help differentiate between vaccines as per the table below. A roll of 500 stickers can be ordered as a standalone product through the CIR inventory.

**Note:** Comirnaty purple cap vaccine will no longer be available in Aotearoa New Zealand from 1 March 2023

## Syringe Labels

**Syringe label for the Comirnaty 15/15 mcg Original/ Omicron Bivalent grey cap vaccine (16+ years if eligible for Booster)**

<p><b>Comirnaty Vaccine</b> Original/Omicron BA.4-5 15/15 mcg dose/0.3mL</p> <p><b>DO NOT DILUTE</b></p> <p><b>Booster dose</b> 16 + years if eligible</p>	Use by DD / MM / YY
	at HH : MM hrs
	Sub batch no. From box
	Box expiry date Sub batch

**Syringe label for the Comirnaty 30 mcg grey cap vaccine (12+ years)**

<p><b>Comirnaty Vaccine</b> 12 + years 30 mcg dose/0.3mL</p> <p><b>DO NOT DILUTE</b></p> <p><b>Primary dose</b></p>	Use by DD / MM / YY
	at HH : MM hrs
	Sub batch no. From box
	Box expiry date Sub batch

**Syringe label for the Comirnaty 10mcg orange cap vaccine (5-11 years)**


<p>Pfizer/BioNTech Comirnaty Vaccine <b>5-11 yrs 10µg / 0.2mL</b></p> <p>Prepared doses used within 6 hours, or by open vial expiry time (whichever is earliest)</p> <p>Vial batch no.</p> <p>Sub batch no. from box</p> <p>Box expiry date Sub batch</p>	Use <b>within 6 hours</b> of draw up
	Use by DD / MM / YY
	at HH : MM hrs
	Prepared by
	at HH : MM hrs
	Checked by

**Syringe label for the Comirnaty 3mcg maroon cap vaccine (6 months – 4 years)**

<p><b>Pfizer/BioNTech Comirnaty Vaccine 6 mths - 4 yrs 3µg/0.2mL</b></p> <p>Prepared doses used within 6 hours, or by open vial expiry time (whichever is earliest)</p> <p>Vial batch no.</p> <p>Sub batch no. From box</p> <p>Box expiry date Sub batch</p>	Use <b>within 6 hours</b> of draw up
	Use by DD / MM / YY
	at HH : MM hrs
	Prepared by
	at HH : MM hrs
	Checked by

<b>Syringe label for the Nuvaxovid vaccine (12+ years)</b>	<b>Novavax COVID-19 Vaccine 0.5mL</b>	
	For IM injection ONLY Prepared doses used <b>within 6 hours</b> or by vial expiry time (whichever is earliest)	<b>Use by</b> DD / MM / YY
	<b>Batch number</b>	<b>at</b> HH : MM hrs
	<b>Vial expiry date</b>	<b>Prepared by</b>
		<b>Checked by</b>

<b>Vaccine packs</b>	
<p><b>Vaccine pack for the Comirnaty Original/Omicron BA.4-5 15/15mcg grey cap vaccine (16+ years if eligible for booster dose)</b></p> <p><b>Note:</b> the sub-batch labels on the vaccine pack for the Comirnaty grey cap vaccine (16+ year olds) is grey to assist with differentiation.</p>	
<p><b>Vaccine pack for the Comirnaty 30mcg grey cap vaccine (12+ years)</b></p> <p><b>Note:</b> the sub-batch labels on the vaccine pack for the Comirnaty grey cap vaccine (12 + year olds) is purple to assist with differentiation.</p>	
<p><b>Vaccine pack for the Comirnaty 10mcg orange cap vaccine (5-11 years)</b></p> <p><b>Note:</b> the sub-batch labels on the vaccine pack for the Comirnaty orange cap vaccine (5–11-year-olds) is orange to assist with differentiation.</p>	

<p><b>Vaccine pack for the Comirnaty 3mcg maroon cap vaccine (6 months - 4 years)</b></p> <p><b>Note:</b> the sub-batch labels on the vaccine pack for the Comirnaty maroon cap vaccine (6 months – 4-year-olds) is maroon to assist with differentiation.</p>	
<p><b>Vaccine pack for the Nuvaxovid vaccine (12+ years)</b></p>	

## 8.3 Cold chain storage

All facilities must hold cold chain accreditation as per the *National Standards for Vaccine Storage and Transportation for Immunisation Providers 2017* (the National Standards). The cold chain accreditation expiry date and back up fridge for each facility must be recorded in the CIR.

Vaccine must be stored and transported in cold chain accredited conditions. NPHS Te Whatu Ora requires any individuals responsible for handling the vaccine to have completed the appropriate cold chain training.

Further information on cold chain management is available in **section 2.1** of the *Immunisation Handbook*. See also the manufacturer’s specifications for approved product handling, available at: <https://www.medsafe.govt.nz/profs/datasheet/c/comirnatyinj.pdf>.

See shelf life of vaccines in the table below. Storage should protect from light.

**Table 8.1 – vaccine shelf life**

	State	At +2°C to+8°C	At ambient temperature
<p><b>Comirnaty 15/15mcg Original/ Omicron BA.4/5 grey cap (16+yrs if eligible for booster)</b></p> <p><b>DO NOT DILUTE</b></p>	Vaccine is prediluted	<p><b>Up to 10 weeks</b> after removal from the ULT freezer.</p> <p><b>Note:</b> Please always follow the ‘in-use’ expiry date label on the vaccine pack. Vaccines can be administered on day of expiry (up to midnight).</p> <p>Please note the expiry date on <b>the vial</b> is only relevant to ULT freezer storage.</p>	<p><b>Unopened vial up to 2 hours</b> (8°C to 30°C)</p> <p><b>Note:</b> there is up to 12 hours allowed, however, 2 hours is a precautionary measure.</p> <p><b>Up to 6 hours when drawn up into syringe or by time written on the vial (whichever is earliest)</b> (up to 30°C)</p>
<p><b>Comirnaty 30 mcg grey cap (12+yrs)</b></p> <p><b>DO NOT DILUTE</b></p>	Vaccine is prediluted	<p><b>Up to 10 weeks</b> after removal from the ULT freezer.</p> <p><b>Note:</b> Please always follow the ‘in-use’ expiry date label on the vaccine pack. Vaccines can be administered on day of expiry (up to midnight).</p> <p>Please note the expiry date on <b>the vial</b> is only relevant to ULT freezer storage.</p>	<p><b>Unopened vial up to 2 hours</b> (8°C to 30°C)</p> <p><b>Note:</b> there is up to 12 hours allowed, however, 2 hours is a precautionary measure.</p> <p><b>Up to 6 hours when drawn up into syringe or by time written on the vial (whichever is earliest)</b> (up to 30°C)</p>
<p><b>Comirnaty 10mcg orange cap (5-11yrs)</b></p>	Undiluted	<p><b>Up to 10 weeks</b> after removal from the ULT freezer.</p> <p><b>Note:</b> Please always follow the ‘in-use’ expiry date label on the vaccine pack. Vaccines can be administered on day of expiry (up to midnight).</p> <p>Please note the expiry date on <b>the vial</b> is only relevant to ULT freezer storage.</p>	<p><b>Up to 2 hours</b> (8°C to 30°C)</p> <p><b>Note:</b> there is up to 12 hours allowed however, 2 hours is a precautionary measure.</p>
	Diluted	<p><b>Up to 12 hours in vial</b> (or until the end of the day it was prepared on)</p>	<p><b>Up to 6 hours when drawn up into syringe or by time written on the vial (whichever is earliest)</b> (up to 30°C)</p>
<p><b>Comirnaty maroon cap (6 months to 4 years)</b></p>	Undiluted	<p><b>Up to 10 weeks</b> after removal from the ULT freezer.</p> <p><b>Note:</b> Please always follow the ‘in-use’ expiry date label on the vaccine pack. Vaccines can be administered on day of expiry (up to midnight).</p> <p>Please note the expiry date on <b>the vial</b> is only relevant to ULT freezer storage.</p>	<p><b>Up to 2 hours</b> (up to 30°C)</p> <p><b>Note:</b> there is up to 12 hours allowed however, keeping it at 2 hours as a precautionary measure.</p>



	Diluted	<b>Up to 12 hours in vial</b> (or until the end of the day it was prepared on)	<b>Up to 6 hours when drawn up into syringe or by time written on the vial (whichever is earliest)</b> (up to 30°C)
<b>Nuvaxovid</b>  <b>DO NOT DILUTE</b>	Unopened	<b>Up to 9 months</b>	<b>Up to 6 hours</b> (up to 25°C)
	Punctured vial	<b>Up to 6 hours</b>	<b>Up to 6 hours</b> (up to 25°C) <b>Up to 6 hours when drawn up into syringe or by time written on the vial (whichever is earliest)</b> (up to 25°C)

### 8.3.1 Process for Refrigeration Failure or Temperature Excursion.

In the event of refrigeration failure which results in a temperature excursion of the vaccine, follow the steps below.

**Table 8.2 – refrigeration failure procedure**

<b>Step 1</b>	
<b>Label the vaccines 'not for use' and in the event:</b>	
<ul style="list-style-type: none"> <li>The refrigerator is currently running within the +2°C to +8°C range, the labelled vaccines are to be retained in your refrigerator.</li> </ul>	<ul style="list-style-type: none"> <li>The refrigerator is not within the +2°C to +8°C range, reversible causes should be considered (door open, power interruption). If no cause found, the labelled vaccines are to be packed into a chilly bin, with a temperature monitoring device and transported to the nearest back-up provider (details for this are in your cold chain policy and in the CIR).</li> </ul>
<b>Step 2</b>	
<b>Contact your local immunisation coordinator for advice and further actions.</b>	
<ul style="list-style-type: none"> <li>Email is monitored from 8.30am to 5.00pm weekdays or contact the Clinical Advice line on <b>0800 IMMUNE (466 863)</b> for guidance up to 8.00pm weekdays or on weekends.</li> </ul>	<ul style="list-style-type: none"> <li>Northern: Lisa Box (<a href="mailto:lisa.box@auckland.ac.nz">lisa.box@auckland.ac.nz</a>)</li> <li>Midland: Karen McKellar (<a href="mailto:Karen.McKellar@auckland.ac.nz">Karen.McKellar@auckland.ac.nz</a>)</li> <li>Central: Abbey Palmer (<a href="mailto:Abbey.Palmer@auckland.ac.nz">Abbey.Palmer@auckland.ac.nz</a>)</li> <li>Southern: Sue Rogers (<a href="mailto:Sue.Rogers@auckland.ac.nz">Sue.Rogers@auckland.ac.nz</a>)</li> </ul>
<b>Step 3</b>	
<b>Document the steps and actions taken.</b>	

## 8.4 Movement of vaccine

Vaccine can be moved around a vaccination facility carefully if required. Avoid any unnecessary movement or handling.

**The vaccine must not be shaken at any stage of transportation, preparation, or administration.**

**Note:** If vials are dropped, or there is another reason for concern about whether the vaccine is still viable, contact **IMAC for advice on 0800 IMMUNE (466 863)**, option 1 (health professionals) and then option 2 (COVID-19 vaccinator support).

## 8.5 Repacking vaccine at Health District facilities

Re-packing only applies to Pfizer vaccines that come in different size packs.

- **Who can re-pack vaccines?**  
Only a Health District hospital pharmacy department can repack the vaccine packs down to distribute to a vaccinator or site. This function is actioned under their hospital pharmacy's Licence to operate Pharmacy and only able to do so for supply within their Health District. In this circumstance, Health District means within the Health District legal entity.
- **Who cannot re-pack vaccines?**  
Health District hospital pharmacy departments are not able to re-pack the vaccine packs for supply to providers outside of their Health District.
- **What if a hospital pharmacy is required to repack the vaccine packs?**  
The Health District hospital pharmacy department will need a packing licence issued to them by Medicines Control.

## 8.6 Transportation of vaccine to other locations

### 8.6.1 Permissible Stock Movement

Sites who have received their vaccine stock from a Health District Pharmacy can contact the pharmacy to organise a stock movement. The Health District Pharmacy can move whole packs, under their wholesale license. Note: all movements must comply with the **National Standards for Vaccine Storage and Transportation for Immunisation Providers 2017**.

This has significant resource implications for the Health District Pharmacy therefore tight stock management is important to minimise waste, and if a stock transfer is necessary, please plan ahead to provide maximum time to support Health District Pharmacy processes.

A provider may take their own vaccine offsite for outreach/home visiting purposes. All cold chain requirements must be met.

No other transportation of vials is permissible.

## 8.6.2 Restrictions on Transport Durations

**For Comirnaty vaccines:**

- **15/15mcg grey cap (age 16+ years if eligible for booster)**
- **30mcg grey cap (age 12+ years)**
  
- **10mcg orange cap (age 5 to 11 years)**

3mcg orange cap (age 6 month to 4 years) There is no limit on the transit time of an unopened vials transported at 2°C to 8°C however, normal shelf-life limits apply.

**For Nuvaxovid (12+ years) vaccine:**

There is no limit on the transit time of an unopened vial of the Nuvaxovid vaccine transported at 2°C to 8°C however, shelf-life limits apply.

## 8.7 Transportation of diluted or drawn-up vaccine

### 8.7.1 Transportation of pre-drawn syringes

The syringes must be appropriately labelled (content, volume, batch, and expiry).

### 8.7.2 Bulk preparation of pre-drawn syringes

The bulk preparation of pre-drawn vaccine to be transported to another location is regarded as compounding and is not permitted unless it is undertaken in an approved facility (such as a hospital pharmacy aseptic unit, or a third-party commercial compounder) with appropriate checks, documentation, and regulator audit.

**Note:** Diluted Pfizer vaccine can be in transit for up to six hours at 2°C to 30°C. Microbiological risks and package integrity, particularly for prepared dosing syringes, are the responsibility of the preparer during transportation of diluted vaccine.

# 9 Vaccine and consumables ordering and delivery

## 9.1 Vaccine ordering

### 9.1.1 Inventory order

Vaccine stock (inventory) can be ordered using the CIR in two ways:

- Direct from the national distribution hubs using a supplier order (see section below),  
or
- From another vaccine site using a **transfer order** (see section below).

See the Standard Operating Procedure (SOP) for order fulfilment at this **SOP for order fulfilment CIR link**

### 9.1.2 Supplier order

This is an order where the stock will come directly from a national distribution hub and the order must be approved by NPHS Te Whatu Ora team. Users must be associated with a location to place a supplier order.

Further details regarding how to log into the CIR can be found in the quick guides, videos and detailed training guide at this **link**.

See the Standard Operating Procedure (SOP) for order fulfilment at this **SOP for order fulfilment CIR link**

#### **Cancelling orders**

Orders can be cancelled before they are approved by NPHS Te Whatu Ora. This is to allow corrections to an order that might be incorrect or orders that are no longer required.

### 9.1.3 Transfer orders

This is a transfer between two locations. It is used routinely to transfer stock between Health District Hospital Pharmacies and mobile vaccination sites. For fixed vaccination sites, the transfer order process is only used for surge/back-up transfers for delivery from Health District Hospital Pharmacies, or end of day returns between two locations. Users must be associated with a location to place a transfer order.

See the Standard operating procedure (SOP) for inventory management on this **SOP for inventory management CIR link**

**Table 9.1 – ordering information required**

Details	New Process
<ul style="list-style-type: none"> <li>• <b>Each site will be allocated a day of the week for delivery. High volume sites may have more than one designated delivery day per week</b></li> <li>• <b>Vaccine orders must be submitted before 10am the day before your allocated delivery day.</b></li> <li>• <b>For the Pfizer Vaccine: a facility should consider the size of the packs they are ordering and their ability to break down packs to avoid unnecessary vaccine movement or wastage.</b></li> </ul>	<ul style="list-style-type: none"> <li>• Vaccine orders must be made through the CIR inventory portal.</li> <li>• The CIR inventory portal will only allow orders for deliveries on the allocated delivery day(s).</li> <li>• If an order is <b>not placed before</b> 10am the day before your allocated delivery day, the Health District will need to submit a request for an 'out-of-cycle' delivery to NPHS Te Whatu Ora CST Logistics Desk.               <ul style="list-style-type: none"> <li>• Orders for Monday delivery are confirmed Thursday/Friday</li> <li>• Orders for Tuesday delivery are confirmed Friday/Monday</li> </ul> </li> </ul>

## 9.1.4 Vaccine delivery schedule

- **How often will I receive vaccine deliveries?**  
The frequency will depend on your typical volume and frequency. For example, a site with higher volumes can receive more regular shipments while lower volume sites or sites only operating on one day a week may choose to receive only one shipment per week.
- **Can my delivery schedule change?**  
The schedule will be discussed and agreed with Health Districts or providers and can be reviewed when required.
- **What if I miss the cut off (by 10am the day before) for ordering vaccines?**  
If you need to order vaccine urgently prior to your next designated delivery day, notify your Health District and they will need to send an 'out-of-cycle' delivery request to the CST Logistics Desk.
- **Where will the vaccine be shipped to?**  
To the location agreed with the Health District or provider.
- **How will I know what vaccines I am due to receive?**  
CIR shows designated delivery days and incoming orders.
- **What if I receive a shipment I am not expecting or don't receive a shipment when I am expecting one?**  
Delivery tracking will be managed centrally by NPHS Te Whatu Ora. Please contact NPHS Te Whatu Ora logistics customer services team.

## 9.1.5 Vaccine Unit sizes and Dimensions

	Unit Size	Unit Dimensions
<b>Comirnaty 15/15 mcg grey cap (16 + years if eligible for booster)</b>	10 multidose vial packs	37mm x 47mm x 89mm
	5 multidose vial pack (available to order TBC)	130mm x 65mm x 45mm
	2 multidose vial pack (available to order TBC)	130mm x 65mm x 45mm
<b>Comirnaty 30 mcg grey cap (12 + years)</b>	10 multidose vial packs	37mm x 47mm x 89mm
	5 multidose vial pack (available to order TBC)	130mm x 65mm x 45mm
	2 multidose vial pack (available to order TBC)	130mm x 65mm x 45mm
<b>Comirnaty 10 mcg orange cap (5-11 years)</b>	10 multidose vial packs	37mm x 47mm x 89mm
	5 multidose vial pack (available to order 27/02/23)	130mm x 65mm x 45mm
	2 multidose vial pack (available to order 27/02/23)	130mm x 65mm x 45mm
<b>Comirnaty 3 mcg maroon cap (6 months – 4 years)</b>	10 multidose vial packs	37mm x 47mm x 89mm
	5 multidose vial pack (available to order 27/02/23)	130mm x 65mm x 45mm
	2 multidose vial pack (available to order 27/02/23)	130mm x 65mm x 45mm
<b>Nuvaxovid (12+ years)</b>	10 multidose vial packs	92mm x 36mm x 62mm

## 9.1.6 Consumables Kits

Please note kits will be discontinued in March 2023 due to all Adult COVID-19 vaccine are now prediluted. Kit items can be ordered individually.

**Table 9.2 – consumable kits**

Item	Material number	Notes	100 Dose Kit	700 Dose Kit
<b>Nipro 25G Standard Needle</b>	1170095	Drawing-up needle for saline – 1 syringe per vial dilution	20 (partial pick)	200 (2 cartons)
<b>BD 3 mL Syringe</b>	1165009	Drawing-up syringe for saline – 1 syringe per vial dilution	20 (partial pick)	100 (1 carton)
<b>Vernacare LDS Needle</b>	1165446	Administering needle – 1 per dose	100 (1 carton)	700 (7 cartons)
<b>Unifix 1 mL Syringe</b>	1169565	Administering syringe – 1 per dose	100 (1 carton)	700 (7 cartons)

**Table 9.3 – order as required**




Item	Purpose	Carton size
<b>Biohazard Yellow Bags</b>	Disposal of waste	50
<b>Sharps Containers – 15 L (5 pack)</b>	Disposal of sharps	5 x 15L
<b>Antiseptic Swabs</b>	Vial disinfectant	200
<b>Non-Woven Swab</b>	Swab	100
<b>Dermaplast Sensitive Injection Plaster</b>	Plaster	250
<b>10 mL Saline (1.8mL per vial)</b>	Diluent	100 x 10mL
<b>SOL-M 1ml Syringe + 25Gx16mm Needle</b>	Administration needle for smaller arms	100
<b>Vernacare LDS Blue Needle 23Gx38mm</b>	Administration needle for larger arms	100
<b>Vernacare LDS Orange Needle 25Gx25mm</b>	Administration needle standard	100
<b>Unifix 1 mL Syringe</b>	Administration syringe	100
<b>BD Flu+ 1mL Syringe + 23Gx25mm LDS Needle</b>	Administration syringe with needle attached	100
<b>Nipro 25G Standard Needle</b>	Drawing-up needle for saline	100
<b>BD 3 mL Syringe</b>	Drawing-up syringe for saline	100

**Table 9.4 – syringe and needle for prediluted vaccines (Nuvaxovid)**

Item	Material number	Carton size
<b>BD Flu Plus 0.1-1mL and 23G 1" LDS needle</b>	1173501	200

## 9.2 Delivery to sites

Figure 9.1 – delivery security

 <b>Warehouse/distribution provider</b>	 <b>Health District facility or vaccination facility</b>	 <b>Vaccination site</b>
<b>Role of NPHS Te Whatu Ora</b>	<b>Role of Health District</b>	<b>Vaccine handover</b>
<p>NPHS Te Whatu Ora will arrange secure transportation of the large quantities of vaccine from the vaccine distribution provider to the cold chain storage facility (such as Health District facility or vaccination site) using a NPHS Te Whatu Ora-contracted courier.</p>	<ul style="list-style-type: none"> <li>• If the vaccine is transported to a Health District cold chain storage facility, secure transportation of the vaccines from that facility to the vaccination sites becomes the responsibility of the relevant Health District or provider.</li> <li>• In the event vaccines are to be transported from a local facility to the vaccination site, the unique circumstances of such transportations should be considered in the site risk assessment.</li> <li>• In the event couriers or authorised personnel (such as vaccinators, administrators, or security) are conducting the transport, NPHS Te Whatu Ora recommends there should be direct travel to the vaccination site (that is, no transit points).</li> </ul>	<p><b>Note:</b> There should be a local procedure in place to ensure the person responsible for transporting the vaccine can be identified. This is to ensure the Health District, or provider has complete confidence they are handing over the vaccine for delivery to the appropriate person. There is no requirement for the person to be a vaccinator.</p>

### Shipper boxes that may be used for transportation from warehouse/distribution provider



*Credo Cube*



*Cool Green Cell*



Please note placement of the all-in-one TrackIT V3 temperature/ tracking device with LCD screen on the inside of the box in vaccine deliveries from DHL (see below):



## 9.2.1 Delivery temperature and expiry dates

Check the sub batch label on the vaccine pack for the expiry date of the vaccine.

## 9.2.2 Vaccine stock/inventory management

- Stock should be used on a **first to expire first out** (FEFO) basis, to ensure waste due to expiry is minimised.
- If there is any concern that a site has excess stock, this should be reported to the Health District who can arrange redistribution.
- Sites should hold two weeks of stock cover.




### Process


Site stock on hand should be managed through the CIR Inventory.




1. Once stock is delivered to a site:
  - Check and verify batch details against details on the order record. Report any discrepancy to the NIP logistics team.
  - Mark stock as receipted in the CIR Inventory once the site has accepted the stock
2. Check the vial and follow in-use expiry on vaccine packs. Due to vaccine expiry extensions, vial expiry may have passed, but the vaccine is still viable.
  - During the preparation of doses and document this on the drawn-up doses label
  - Before administration of the vaccine
  - At the end of the day check stock
3. Discard any expired vaccines and record this as waste in the CIR Inventory (see section '**Recording vaccine waste**').
4. Any consumption and wastage must be recorded in the CIR Inventory daily.
5. Once consumption is recorded in the CIR Inventory, all remaining stock on site must be checked against the stock showing in the CIR Inventory to ensure that there are no discrepancies.
6. Any discrepancies must be investigated and captured in the CIR Inventory as stock adjustment.

For more detail see the Standard operating procedure (SOP) for inventory management on this [SOP for inventory management CIR link](#)

**Table 9.4 – site delivery and receipt process**

Step	Action
 <p><b>Health District/provider logistics lead provides site contact and delivery details</b></p>	<p><b>Site checklist</b></p> <p>The site checklist must be completed prior to the site commencing vaccinations (see <b>Appendix A</b>).</p> <p><b>Site contact</b></p> <ul style="list-style-type: none"> <li>The Health District or provider logistics lead must provide NPHS Te Whatu Ora with: <ul style="list-style-type: none"> <li>a site contact (a named role and a phone/mobile number)</li> <li>detailed delivery instructions, including address and any special instructions (such as separate entrances and so on).</li> </ul> </li> <li>Submit this information using the New facility/ site set-up form (<b>Appendix B</b>) <i>at least 5 days prior</i> to ordering vaccines for that site.</li> </ul> <p><b>Availability of site contact</b></p> <ul style="list-style-type: none"> <li>The site contact should be regularly available on site to accept deliveries. This will minimise the administration involved changing the site contact person, for example.</li> <li>Please notify <b>urgent</b> site contact changes to NPHS Te Whatu Ora logistics team.</li> </ul> <p><b>Cold chain accreditation</b></p> <ul style="list-style-type: none"> <li>NPHS Te Whatu Ora recommends individuals handling vaccines are cold chain accredited; however, this is not a requirement.</li> </ul>
 <p><b>Vaccine distribution provider packs and ships vaccine</b></p>	<p><b>Ship under cold chain conditions</b></p> <ul style="list-style-type: none"> <li>The vaccine distribution provider will pack and ship the vaccine under cold chain conditions in shipping boxes, depending on delivery destination, at +2°C to +8°C.</li> </ul>
 <p><b>Site contact receives the package</b></p>	<p>The courier will hand the package to the site contact. Before signing for the package, the site contact will:</p> <ul style="list-style-type: none"> <li>Confirm the shipping box is addressed to them/their site</li> <li>Provide their identification to the courier for the courier's confirmation</li> <li>Conduct a check of the order immediately while the courier is present (see below)</li> </ul>

Step	Action
<p><b>Site contact checks the temperature datalogger</b></p> <p><b>Econolog Temperature Reading (same device comes in different colours)</b></p>  <p><b>If the temperature datalogger shows:</b></p> <ul style="list-style-type: none"> <li>• <b>Green light flashing once every 10 seconds</b> <ul style="list-style-type: none"> <li>○ <b>The temperature has remained within limits</b></li> </ul> </li> <li>• <b>Red light flashing once every 10 seconds</b> <ul style="list-style-type: none"> <li>○ <b>Excursion has occurred</b></li> </ul> </li> </ul> <p><b>TrackIT V3 Temperature Reading</b></p>  <p><b>If the temperature datalogger shows:</b></p> <p><b>Green light and ✓</b></p> <ul style="list-style-type: none"> <li>• <b>The temperature has remained within limits</b></li> </ul> <p><b>Red light and X</b></p> <ul style="list-style-type: none"> <li>• <b>Excursion has occurred</b></li> <li>• <b>If the screen is not displaying press the button once</b></li> </ul>	<p><b>Check for a temperature excursion</b></p> <p>The site contact must follow the process below:</p> <ul style="list-style-type: none"> <li>• Read the temperature datalogger immediately after opening the box (before removing the vials).</li> <li>• Do not attempt to stop the temperature datalogger</li> <li>• Leave the datalogger in the shipment container.</li> </ul> <p><b>In the event of a temperature excursion</b></p> <ul style="list-style-type: none"> <li>• Quarantine the shipment in cold chain conditions.</li> <li>• Return the shipment container with temperature datalogger inside to the courier.</li> <li>• Contact the NIP Logistics Customer Support Service Team immediately.</li> </ul> <p><b>Temperature excursion – next steps</b></p> <p>The NIP logistics team will advise the site contact on the next steps, such as the need to re-order and use of quarantined vaccines once the temperature report has been reviewed.</p>
 <p><b>Site contact conducts visual check</b></p>	<p><b>Visual check</b></p> <ul style="list-style-type: none"> <li>• The site contact will open the shipping box and the internal vaccine packaging and conduct a visual check of the outer packaging to check for damage and/or leakage. If there is no damage store directly in the fridge.</li> <li>• Each site should check the packing slip to make sure all vaccines have been received</li> </ul>

Step	Action
	<ul style="list-style-type: none"> <li>• If there are any signs of damage to the outer container, inspect the vials inside the package: <ul style="list-style-type: none"> <li>• Broken vials or waste needs to be recorded in the CIR logistics module, but only to the unopened vial stage</li> <li>• Vaccine wasted in opened vials is not required to be recorded in the CIR logistics module.</li> <li>• Please see the <b>Standard Operating Procedures</b> in the <b>Inventory orders</b> section regarding how to record vial consumption and waste.</li> </ul> </li> </ul>
 <p><b>Site contact signs for vaccine package</b></p>	<p><b>Vials intact</b></p> <ul style="list-style-type: none"> <li>• Where the vials are intact and there are no concerns, the site contact will sign for the package.</li> </ul>
 <p><b>Site contact stores vaccine in cold chain accredited conditions</b></p>	<p><b>Store vaccine</b></p> <p>The site contact will then store the vaccine at cold chain conditions in the internal packaging carton it arrived in (not the shipping box (<i>Credo Cube/Cool Green Cell</i>)), but the vaccine pack until the expiry date and time marked on the vaccine pack is reached. Any vials no longer viable must be disposed of following the disposal process detailed below.</p>
 <p><b>Receipting orders</b></p>	<p>When a vaccine or consumables order is received, it must be receipted into the CIR. This enables the movement of the stock from in transit to available for use in the stock on hand.</p> <p>Further details regarding how to log into the CIR can be found in the quick guides, videos, and detailed training guide at <a href="https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Inventory-Orders-Portal-Quick-Step-Guide">https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Inventory-Orders-Portal-Quick-Step-Guide</a>.</p> <p>See the Standard Operating Procedure (SOP) for order fulfilment at this <b>SOP for order fulfilment CIR link</b></p>

## Equipment returns

**Table 9.5 – temperature monitoring, *Shipping boxes* equipment return**

Details	Process
<p><b>Shipping boxes and temperature monitoring equipment should be returned in a timely manner – preferably on the same day as receipt – to ensure there are no interruptions of subsequent vaccine deliveries.</b></p>	<ul style="list-style-type: none"> <li>• Pre-paid stickers will be included with the delivery for returns.</li> <li>• The number on the instructions should be called to arrange collection.</li> <li>• Any fault or damage to the packaging equipment should be reported at the time of return.</li> <li>• <b>Note:</b> Ensure correct removal or crossing-out of the original courier label and original address details to avoid any confusion.</li> </ul>

**Table 9.6 – daily reporting information required**

Vaccination events	Significant events	Stock on hand / stock movements
<ul style="list-style-type: none"> <li>• Sites must ensure vaccination events are recorded in CIR at the time of administration. This enables accurate data for operational reports, such as number of vaccinations completed and other trends.</li> </ul>	<ul style="list-style-type: none"> <li>• Providers need to report significant events on sites such as a significant adverse reaction, or a protest to the Ministry daily as required.</li> </ul>	<p>Providers must ensure the following information is recorded in the CIR inventory portal daily as required:</p> <ul style="list-style-type: none"> <li>• Facility stock on hand</li> <li>• Stock movements from facility to facility</li> <li>• Stock movements from facility to site</li> </ul>

**Table 9.7 – asset management recommended practice**

Recommended practice	Details
<p><b>Collation of site inventory and operations</b></p>	<p>Health Districts or providers may wish to collate daily reporting back from sites on inventory and/or operations to aid the supply of information back to NPHS Te Whatu Ora.</p>
<p><b>Demand planning</b></p>	<p>Maintain a 4-week forward demand plan.</p>
<p><b>Continuous process improvement</b></p>	<p>NPHS Te Whatu Ora welcomes feedback on the immunisation process or recommendations for operational improvements.</p> <p>Please contact your regional liaison to pass on your feedback</p>

## 9.3 Reports available to Health Districts

- **What information is available in reports?**

As the COVID-19 vaccine reporting is linked to the NHI database, requesting any existing NHI data fields (such as ethnicity) to track vaccination rates and meet other reporting needs is valid.

- **How do I request reports?**

Contact your Health District or provider reporting team, who will then submit your request to NPHS Te Whatu Ora's reporting team.

- **Where can I find my reports?**

Once the report is prepared, it will be available in CIR as both a dashboard and downloadable report.

- **Will my reports be refreshed?**

Reports will be updated in real-time.

- **What if I want reports for multiple Health Districts?**

Please specify this at the time of requesting the report.

## 9.4 Vaccine and consumables assets and asset management

An asset is an instance of vaccine stock and vaccine consumables, such as: five pack of vaccine, 15 pack of vaccine, 195 pack of vaccine, or consumable kit.

Assets at a location can be updated through:

- Stock re-work
- Stock adjustment
- Quarantine stock
- Recording consumption, or
- Stock on hand.

Further details regarding how to log into the CIR can be found in the quick guides, videos, and detailed training guide at <https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Inventory-Orders-Portal-Quick-Step-Guide>.

See the Standard operating procedure (SOP) for inventory management at this [SOP for inventory management CIR link](#)

### Recording consumption

It is important to record the consumption of vaccine stock and consumables as stock in consumed or, as a minimum, as part of the daily stocktake. The purpose of this is to give an accurate local, regional, and national view of vaccine stock on hand.

Consumption can be recorded in two ways:

1. Consumption – entering directly what has been consumed
2. Stock on hand – entering a physical count of the stock on hand as part of the daily stock take

Further details regarding how to log into the CIR can be found in the quick guides, videos, and detailed training guide at <https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Inventory-Orders-Portal-Quick-Step-Guide>.

See the Standard operating procedure (SOP) for inventory management at this **SOP for inventory management CIR link**

## Recording vaccine waste

It is important for vaccine sites to record vaccine waste in the CIR Logistics Portal, but only to the unopened vial level (the recording of vaccine wasted at the opened vial level is yet to be determined). This is so that waste can be tracked at a local, regional, or national level.

Further details regarding how to log into the CIR can be found in the quick guides, videos, and detailed training guide at <https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Inventory-Orders-Portal-Quick-Step-Guide>.

See the Standard operating procedure (SOP) for inventory management at this **SOP for inventory management CIR link**

# 10 Disposal of consumables, vaccine, and vaccine packaging

Vaccine disposal and other inventory management topics (outlined below) are available as eLearning modules.

## 10.1 Disposal of consumables

Health Districts and providers are responsible for the disposal of consumables. Consumables should be disposed of according to existing procedures (such as disposal into sharps bin and/or biohazard bags). Local procedures are to be followed to arrange collection of the sharps bin and other medical waste.

## 10.2 Disposal of damaged, empty, and expired vaccine vials

When a possible cold chain breach occurs providers must contact their immunisation coordinator before disposing of any vaccines as per the **National Standards for Vaccine Storage and Transportation for Immunisation Providers 2017**.

The process for destruction and disposal of expired vials is as follows:

1. Remove the lid
2. Deface the vial
3. Place the vial(s) in the Interwaste vial disposal bin
4. Record the wastage in the CIR Inventory



As part of site preparations, Interwaste must be contacted at least 5 business days in advance of your site going live to request a vial disposal bin to be delivered to the site. Contact Interwaste on 0800 102 131 (their call centre is available from 8am-5pm weekdays). For more information see the **Ordering Interwaste vial disposal bin** section above.

Interwaste will provide a 20-litre sized container in which to dispose expired (full), empty, broken, or damaged vials. Please note, expired vials should be defaced before disposal. When the container is almost full, contact Interwaste on 0800 102 131 to arrange its pick-up. Interwaste will deliver a new disposal container at the same time they remove the existing container. Interwaste will destroy the vials in an appropriate manner.

Ensure the lid of the Interwaste disposal container remains closed when not in use.



Figure 10.1 – disposal bin

## 10.3 Disposal of vaccines drawn up but not administered and empty vaccine syringes

Vaccine doses that have been drawn up but not administered must be disposed of in the sharps bin provided. Similarly, empty/used vaccine syringes should be disposed of in the sharps bin. Seal and remove sharps bins when filled and stored in a secure area for transportation and final disposal

Manage sharps waste as per NZS 4304:2002 *Management of Healthcare Waste*.

## 10.4 Vaccine packaging disposal

Ensure all packaging that the vaccine is sent in is appropriately destroyed to ensure packages cannot be replicated.

Once all vials in a vaccine pack have been used, black-out all vaccine-related information on the label, using a permanent marker. The vaccine pack must be securely destroyed. It can be disposed of in a secure document destruction bin if one is available or a biohazard bag. Packaging must not be disposed of in household waste collection or recycling centres. If additional biohazard bags are required, please advise NPHS Te Whatu Ora logistics when placing the next consumables order.

# 11 Site readiness and closure

## 11.1 Site setup form and site checklist

Complete the site checklists included in **Appendix A** to assess whether the vaccination site is ready to commence vaccinations. Site checklists, upon completion, must be signed by the Health District or provider chief executive, or their delegate, to approve the site is ready. The checklist is then submitted to either the regional account manager or NPHS Te Whatu Ora's logistics team. Primary care providers may be asked to submit site checklists to their Health District rather than NPHS Te Whatu Ora directly.

The new facility/ site set up form (v1.4) form (see **Appendix B**) must be submitted **at least five days prior** to the site commencing vaccinations. This information is used to set up the facility or site in CIR and ensure deliveries are made to the correct address. Care is required to provide accurate information on this form.

## 11.2 Completing a dry run

NPHS Te Whatu Ora recommends a site trial or dry run before beginning vaccinations on site to ensure staff are familiar with their roles and consumer flow can be tested. NPHS Te Whatu Ora's logistics team do not provide dry run packs however, an optional order of consumables can be ordered from NPHS Te Whatu Ora's logistics team which can be used to complete a dry run.

## 11.3 Facility/site closure form

Complete the **facility/site closure form** (see **Appendix C**) as a part of the site and facility closure protocol, and to assess and return stock.

A stocktake of all consumables relating to the COVID-19 Vaccination Rollout must be completed upon site/facility closure. Submit the completed facility and site closure form to NPHS Te Whatu Ora's logistics team and your Health District logistics Lead. This should be submitted a week before the closure or as soon as the closure of the location is known.

## 11.4 Facility moving location

**Facilities** are where vaccines are shipped to, stored, and subsequently distributed to sites. **Sites** are where vaccines are administered.

Facility Moving Location	New Facility Set-up Required
<p>If a facility needs to move their physical location, and there will be continuity of the Cold Chain Accreditation (CCA), the facility can complete part four of the <b>Facility/Site Set-up form</b> (see Appendix B) confirming Health District approval for the move and that CCA will be in place for the new facility.</p>	<p>If Cold Chain Accreditation (CCA) cannot be transferred to the new location and requires reassessment, please:</p> <ul style="list-style-type: none"><li>• Complete the <b>Facility/Site Closure Form</b> and associated procedures.</li><li>• Complete the new <b>Facility/Site Set-up form</b> and <b>associated procedures</b> which will allow the facility to be set up in the CIR for delivery to the new address.</li><li>• Ensure that CCA for the new location has been assessed and approved.</li></ul>

- Cold chain must be maintained during a facility move
- The only circumstance where vaccines can be delivered to an alternative address to that already set up in the CIR, is where it is delivered to the nominated address for cold chain fridge back-up (recorded in the CIR).

# 12 Becoming a COVID-19 Vaccination site

## 12.1 Onboarding

Becoming a COVID-19 vaccination site can be complex, involving engagement with both your local Health District and/or PHO and NPHS Te Whatu Ora. To ensure consumer safety, vaccination sites will need an appointed Clinical Site Lead to navigate the onboarding process. The Clinical Site Lead is accountable for meeting **clinical safety and quality standards** at their site, as well as supporting **planning, clinical governance, quality, and safety management** processes.

Primary Care providers are a critical component of the New Zealand COVID-19 vaccination rollout.

## 12.2 Additional resources

The following supporting documents can be found on the Ministry's website:

- **Technology, User Roles, and Training Matrix**
- **User Onboarding Journey for Book My Vaccine** (also known as NIBS)
- **User Onboarding Journey for COVID-19 Immunisation Register (CIR)**

# Section B: Pathway to COVID-19 vaccination

## Section B: Pathway to COVID-19 vaccination - summary of changes

Version	Date	Section	Summary of Changes
51.0	28/02/23	17	Updated guidance for discontinued Comirnaty 30 mcg purple cap vaccine
		18-19	New sections with key safety points for Comirnaty 30 mcg grey cap and Comirnaty 15/15 mcg Original/Omicron BA.4/5 grey cap
		23	New section 'Preparation of doses' which summarises information previously found in each vaccine section.
		24.3	Removed 'my vaccine pass'

## Section guidance

This section provides operational guidance on the vaccination pathway COVID-19 vaccines, from booking and scheduling to vaccine preparation onto vaccine administration and observation. The first line vaccines where there are no contraindications is the Pfizer-BioNTech Comirnaty vaccines.

## Purpose

The purpose of this section is guiding the vaccinating workforce to *do the right thing* and have the right resources and information available to provide a safe quality vaccination journey for every consumer. It is designed to be applicable to all sites delivering the COVID-19 vaccine and provide guidance and assistance to providers, to maintain public safety and ensure consistent and equitable vaccination practices are in place across New Zealand/Aotearoa.

This section should be read and interpreted alongside the **COVID-19 immunisation policy, Immunisation Handbook 2020**, the Standards, and **IMAC resources**.

### Appendices relevant to this section

- **Appendix G: Vaccination site screening questions**
- **Appendix H: Supported decision-making process**
- **Appendix I: Serious Adverse Event Process** (process steps, SAC examples, notification form)

# 13 Booking and scheduling

The National Immunisation Booking System known as *Book My Vaccine* (BMV) supports a national-led approach to immunising New Zealand/Aotearoa against COVID-19. *Book My Vaccine* supports vaccination sites down to Community Hub level. Use by primary care sites is optional where they only service their own enrolled populations.

For more information, see **Section C: Additional Programme Guidance, Variations, and Incidents**.

Ensure that the scheduling of vaccination appointments avoid over-crowding and allow for physical distancing and other IPC measures. Also, limit the number of accompanying people to only those who need assistance, whether physical or psychosocial.

## 13.1 Booking second doses

### Do not vaccinate less than 21 days

- **The administration of a COVID-19 vaccine second dose at an interval of less than 21 days is not approved by Medsafe and is considered off-label use and must be reported to CARM.**
- **New bookings made through [bookmyvaccine.nz](https://bookmyvaccine.nz) and the COVID-19 vaccine Whakarongorau Aotearoa 0800 28 29 26 is set to three weeks between the two doses.**
- **If consumers have existing vaccination bookings, they can keep their second appointment as it is, or choose to change it. Either way the important thing is that consumers receive two doses of the vaccine to be fully vaccinated.**
- **Consumers should select the appropriate age range when making an appointment**
- **Second doses can be booked for any time after day 21.**
- **The administration of a COVID-19 vaccine at an interval of less than 21 days is not approved by Medsafe and is considered off label use and must be reported to CARM. See Appendix I for more information on incidents.**

**Note:** A prescription from an authorised prescriber is required when using Nuvaxovid as a second primary dose following a non-Nuvaxovid COVID-19 vaccine for a first primary dose, in accordance with Section 25 of The Medicines Act 1981, as it is considered off-label use. This must be documented clearly including the rationale and the informed consent process. A CARM report does not need to be completed if the vaccine has been prescribed by an authorised prescriber.

For more information on dose intervals please see the **COVID-19 Immunisation policy statement on the Ministry of Health website**.

## Administering leftover vaccines

To minimise wastage, the Programme recommends the preparation of a back-up/stand-by list of consumers aligning to the sequencing framework. Leftover diluted and/or drawn vaccine unused at the end of the shift that would expire before the next clinic, may be administered to consumers on the back-up/stand-by list.

The Programme does not require visibility of the back-up/stand-by list; use best judgement to manage this list as to align with the sequencing framework.

# 14 Protecting security and privacy

The vaccination process requires personal, identifying information be collected. In the health sector, NHIs are considered identifiable information as well as standard identifiers such as name, address, and date of birth.

Protecting and treating sensitive health information with respect is important.

- All medical records (such as written consent forms) at vaccination sites are required to be securely stored out of the sight (for example, in a drawer).
  - It is preferable this storage area is locked, or in the constant presence of an authorised person, such as an administrator, a security guard, or a vaccinator.
- At the conclusion of the vaccination event, the Programme recommends that the personal information documentation is taken directly (that is, no transit points) by an authorised person (such as an administrator, a security guard, or a vaccinator) to the site where the record will be held.

In addition to ensuring the security of health records as per above, the following security and privacy factors should be considered:

- Informing consumers why their information is being collected and what it will be used for (for example, that it will not be used for immigration or law-enforcement purposes)
- Consider who may be able to see computer screens that are likely to be used to input personal information
- Ensure passwords and log-in details are kept confidential
- In the event of a likely security or privacy breach advise the relevant Health District or provider privacy officer or contact the Programme's Privacy team as soon as possible
- Securely dispose unnecessary duplicate information
- Ensure confidential conversations occur away from areas where other consumers or members of the public might also access.
- Ensure staff accessing consumer data have completed the appropriate privacy training (e.g., see the **Privacy Commissioner courses link**).

**Note:** Use secure methods when transferring information outside of the core vaccine systems such as USB encryption or accredited online services. Data should be password protected.



# 15 COVID-19 vaccines operational phase

- Use a daily checklist to monitor and ensure IPC and other safety measures are adhered to.
- Consider a daily 'huddle' to enhance teamwork and to highlight any IPC issues.
- Screen all staff for signs and symptoms of COVID-19 at the start of each shift.
- Screen all people arriving for vaccination for COVID signs and symptoms. For additional screening questions see **Appendix G**.
- Ensure the scheduling of vaccination appointments avoids over-crowding and allows for physical distancing and other IPC measures. Also, limit the number of accompanying people to only those who need assistance, whether physical or psychosocial.
- Ensure the appropriate processes are in place to prevent under-age vaccinations – **this is a never event**.
- Ensure the appropriate processes are in place to prevent second dose vaccinations earlier than 21 days – **this is a never event**.
- **Ensure the appropriate processes are in place to ensure consumers are receiving the age-appropriate vaccine.**

**Note:** In the rare occurrence where an authorised prescriber deems the vaccine clinically indicated for a consumer, the authorised prescriber can prescribe the vaccine as off label/unapproved use. This must be documented clearly including the rationale for early second dose and the informed consent process. A CARM report does not need to be completed if the vaccine has been prescribed by an authorised prescriber. Written consent is advised.

## Key IPC measures to implement

Prepare each injection in a clean, designated area.

### Hand hygiene

- At the start of the shift, all vaccination team members are required to wash their hands thoroughly with soap and water and dry them thoroughly or use hand sanitiser.
- Facilitate attending consumers' hand hygiene (as above).
- Vaccinators should perform hand hygiene before putting on and removing PPE, before preparing the vaccine, and between each vaccine administration, preferably using alcohol-based hand sanitisers.
- Gloves are not required and, if used, do not replace the need for hand hygiene between each vaccine administration and for other indications. The use of alcohol hand sanitisers on gloves is strongly discouraged.

### PPE

- PPE is to be selected based on risk assessment as a part of standard precautions.
- In the context of the COVID-19 pandemic, vaccinators should wear PPE appropriate to the area they are working in. For more information visit the **Te Whatu Ora website** .

### Preparation and administration IPC

- Sterile, single use syringes and needles should be used. These should only be removed from their packaging immediately before use.
- Perform hand hygiene before preparing vaccine for delivery
- Prevent contamination of the vials by wiping the access diaphragm (septum) with 70% alcohol (isopropyl alcohol or ethanol) on a swab or cotton wool ball before piercing the vial and allow to air dry. If the top of the vial is accidentally touched during drawing up it must be re-wiped (repeat this step).
- Adhere to IMAC guidance for the drawing up of vaccine and skin preparation at the site of injection.
- Discard used syringes and needles as a single unit into a sharps container immediately after administering the vaccine

# 16 Obtaining informed consent

Prior to administering the vaccination, the registered health professional must obtain informed consent, per the *Code of Health and Disability Services Consumers' Rights* (the Code). The steps to recording the outcome of the informed consent question is:

- The vaccinator or an administrative support person must record in CIR the consumer's consent to approve or decline the administration of the vaccine.
- The Programme assumes verbal consent is agreeable in most situations.
- Written consent can be considered in the following situations below:
  - a. where there are significant risk of adverse effects to the consumer, per **clause 7(6c) of the Code**
  - b. if it is being prescribed. For more information, please refer to the below 'Prescription' section.
  - c. if this is the provider's or vaccinator's preference, for example, in aged residential care settings.
- Where written consent is recorded under points a. b. and/or c. above, the forms do not need to be uploaded to CIR; rather, the provider is responsible for ensuring the forms are archived as a part of that consumer's clinical record.
- If written consent forms are unable to be archived in the consumer's clinical record, then this must be uploaded onto CIR. Once this is complete the record can be destroyed.

Where a consumer is not competent to make an informed choice and give consent for their vaccine, someone who has the legal right can make decisions on the consumer's behalf; namely a legal guardian or someone who currently holds Enduring Power of Attorney for personal care and welfare.

See **Appendix H** which displays the process for consumers requiring support to consent to the COVID-19 Vaccination. Any supported decision-making conversations should be documented in the notes section of CIR. For more information regarding obtaining informed consent, see the *Immunisation Handbook, chapter 2*.

For more information regarding supported decision making, or to access the training module specific to COVID-19 Vaccine Supported Decision Making, see IMAC Learning Courses at **IMAC Learning**.

## Obtaining written consent for the Nuvaxovid vaccine

The Programme requires written consent to be obtained before administering the Nuvaxovid vaccine as a second primary dose after a non-Nuvaxovid vaccination.

## Informed consent for consumers aged 12 to 15 years

Under the code of rights, every consumer, including a child, has the right to the information they need to make an informed choice or to give informed consent. Therefore, a young person aged 12-15 years can provide their own informed consent or refusal to consent if they are deemed competent to give consent, and a parent or guardian does not need to provide consent or be present. Some of these young people may choose to have their parent or guardian consent on their behalf and that is fine.

### Verbal or written consent for consumers aged 12 to 15 years

Informed consent for consumers aged 12-15 years can be verbal. However, written consent can be required if it is the provider's or vaccinator's preference, and like with all consumers, must be obtained if there is significant risk of adverse effects.

## 16.1 Prescription

A prescription from an authorised prescriber is required when a vaccine is being administered off-label under **Section 25 of The Medicines Act 1981**, such as when a Medsafe approved medicine is being used for an un-approved use. However, no prescription from an authorised provider is required if the administration is authorised under **section 34A of The Medicines Act 1981** which empowers the Director-General of Health to authorise, by Notice, the use of a consented COVID-19 vaccine otherwise than in accordance with the approved data sheet.

For the list of authorised prescribers please refer to the **Medsafe website**.

*When a prescription is used, it is recommended that written consent is completed. In this instance it means that the prescriber completes and signs the written consent form. However, if the prescriber is not available to sign the written consent form, the Clinical Lead can complete the form. The prescription and written consent form can be uploaded in the CIR.*

### 16.1.1 Written consent forms

Written consent forms must be managed on-site or by a centralised administration team. Given the information on the written form contains personal information, **forms must be always held and transported securely** (for example, in a locked cabinet/drawer, a tracked courier bag, or other secure container when transported between locations). The consumer may also decide to take the written consent form with them.

If providers choose to upload written consent forms the person uploading, for example the administrator, must scan each form to their computer, locate the consumer's CIR record, then upload the scanned form/s to the consumer's CIR record; delete the local copy and securely destroy the written form. When necessary, the written form may be kept for a few days or weeks to check for inaccuracies in transcribing before the written forms are destroyed.

**Note:** Instructions for uploading files to CIR are included in the CIR eLearning module.

## 16.1.2 Variations to consent forms

As of 6 May 2021, there is now only one NPHS Te Whatu Ora consent form (the Group 1a version m has been withdrawn). The COVID-19 consent form is available via the National Immunisation Programme's Dropbox and the Ministry of Health website. Please ensure you are using the latest version.

## 16.1.3 Vaccine safety and additional considerations for consumers aged 12 to 15 years

Similarly, as with consumers over the age of 16 years, it is important to assess the administration site and select the correct needle length. Most commonly, the same needles used for adults would be used for consumers aged 12-15 years.

# 17 Comirnaty purple cap COVID-19 vaccine (for ages 12 years and over)



As of 1 March 2023, Comirnaty purple cap will no longer be available in Aotearoa New Zealand.

At end of day 28 February 2023 all Comirnaty purple cap vials will be disposed in Interwaste vial disposal bins and recorded as wastage in CIR inventory.

Comirnaty 30 mcg grey cap will be available for the primary course for those 12 years and over. See section 18 for operational guidance.

Comirnaty 15/15 mcg Original/Omicron BA.4/BA.5 will be available for booster doses for those 16 years and over.

# 18 Comirnaty 15/15mcg Original/ Omicron BA.4/5 grey cap COVID-19 vaccine (for ages 16 years and over)

The key safety points are:



- Approved for use for consumers aged 16 years or over receiving a booster dose
- The Comirnaty 15/15mcg Original/ Omicron BA.4/5 grey cap COVID-19 vaccine **does not need** to be diluted
- There are 6 doses per vial
- For all vaccinator resources and materials related to Comirnaty 15/15mcg vaccine please refer to the **IMAC website**.

This is the first line booster vaccine where there are no contraindications to the Pfizer-BioNTech Comirnaty vaccine.

A prescription from an authorised prescriber is required when using the Comirnaty 15/15mcg Original/ Omicron BA.4/5 grey cap vaccine as a primary course dose, in accordance with **Section 25 of The Medicines Act 1981**, as it is considered off-label use. Written consent is required for all consumers receiving an 'off label' dose of the Comirnaty 15/15 mcg Original/ Omicron BA.4/5 grey cap vaccine.

# 19 Comirnaty 30mcg grey cap COVID-19 vaccine (for ages 12 years and over)

The key safety points are:



- Approved for use for consumers aged 12 years or over receiving their **primary course**
- The Comirnaty 30 mcg grey cap COVID-19 vaccine **does not need** to be diluted
- There are 6 doses per vial
- To be considered as a booster dose at less than 6 months after a previous dose, then written consent and a prescription are required.
- This is the first line vaccine where there are no contraindications to the Pfizer-BioNTech Comirnaty vaccine.
- For all vaccinator resources and materials related to Comirnaty 30mcg vaccine please refer to the **IMAC website**.

A prescription from an authorised prescriber is required when using the Comirnaty 30 mcg grey cap vaccine as a booster dose administered at less than 6 months from a previous dose, in accordance with **Section 25 of The Medicines Act 1981**, as it is considered off-label use. Written consent is required for all consumers receiving an 'off label' dose of the Comirnaty 30 mcg grey cap vaccine.

## 20 Comirnaty 10mcg orange cap COVID-19 vaccine (for ages 5 to 11 years)

The key safety points are:



- Approved for use for children/tamariki aged 5 to 11 years
- The Comirnaty orange cap vaccine dose is different to the Comirnaty purple cap dose (12+ years) and the Comirnaty maroon cap dose (6 months to 4 years)
- There are 10 doses per vial
- If the consumer receives the Comirnaty orange cap dose (for ages 5-11) and then turns 12 before their second dose, they will receive the adult dose of the Comirnaty vaccine for subsequent doses.
- For all vaccinator resources and materials related to Comirnaty 10mcg vaccine please refer to the **IMAC website**.



## 20.1 Site readiness

If sites are new to vaccinating the Comirnaty orange cap vaccines (for ages 5 to 11 years) it is recommended a Comirnaty orange cap site, check list is completed

Comirnaty orange cap Site Checklist	Y / N
Site Workforce Police safety check are up to date	Y <input type="checkbox"/> N <input type="checkbox"/>
Vaccinators administering the Comirnaty orange cap vaccine must complete the Paediatric COVID-19 Vaccinator Education Course (IMAC link)	Y <input type="checkbox"/> N <input type="checkbox"/>
Child safe Environment	Y <input type="checkbox"/> N <input type="checkbox"/>
SOP preparation of Comirnaty orange cap doses	Y <input type="checkbox"/> N <input type="checkbox"/>
Child friendly resources (distraction posters can be found on the IMAC website)	Y <input type="checkbox"/> N <input type="checkbox"/>
Child-suitable bag valve mask (BVM or 'ambu bag') resuscitator is required, airways (optional) and any other emergency equipment to respond to a serious adverse event. <b>Note:</b> See A4.6. Minimum staff and equipment requirements for vaccination services in <b>Appendix 4</b> of the Immunisation Handbook (2020)	Y <input type="checkbox"/> N <input type="checkbox"/>
Consumer collateral	Y <input type="checkbox"/> N <input type="checkbox"/>
Dry Run	Y <input type="checkbox"/> N <input type="checkbox"/>
Wet Run	Y <input type="checkbox"/> N <input type="checkbox"/>

## 20.2 Vaccine safety and additional considerations for consumers aged 5 to 11 years

With consumers the age of 5 to 11 years, it is important to use the correct needle length. For children/tamariki under the age of 7 years a 16 mm length needle should be used. For children/tamariki ages 7 to 11 years clinical judgement should be used to determine if a longer needle is required (25mm). Use of a shorter needle risks delivering the vaccine subcutaneously as opposed to intramuscularly, which has the potential to underdose. For more information on needle length, refer to the *Immunisation Handbook*.

### Ensuring young people have adequate understanding of the vaccine and can provide informed consent

Training and guidance material are available to support vaccinators to gauge consumer's ability to provide informed consent. It is important that a robust conversation occurs prior

to vaccination, where the child or their parent/ legal guardian/ enduring power of attorney has an opportunity to have any questions answered and concerns addressed.

## 21 Comirnaty 3mcg maroon cap COVID-19 vaccine (for ages 6 months to 4 years)

The key safety points are:



- Approved for use for children/tamariki aged 6 months to 4 years
- The Comirnaty maroon cap vaccine dose is different to the Comirnaty purple cap dose (12+ years) and the Comirnaty orange cap dose (5 to 11 years)
- There are 10 doses per vial
- If the consumer receives the Comirnaty maroon cap dose (for ages 6 months -4 years) and then turns 5 before their second or third dose, they will receive Comirnaty orange cap (5 to 11 years) for subsequent doses.
- For all vaccinator resources and materials related to Comirnaty 3mcg vaccine please refer to the **IMAC website**.

### 21.1 Vaccine safety and additional considerations for consumers aged 6 months to 4 years

With consumers the age of 6 months to 4 years, it is important to use the correct needle length for the child being vaccinated as well as the area of their body the vaccine is to be given into (ie deltoid vs vastus lateralis). For more information on needle length, refer to the *Immunisation Handbook*.

## Ensuring young people have adequate understanding of the vaccine and can provide informed consent

Training and guidance material are available to support vaccinators to gauge consumer's ability to provide informed consent. It is important that a robust conversation occurs prior to vaccination, where the child or their parent/ legal guardian/ enduring power of attorney has an opportunity to have any questions answered and concerns addressed.

# 22 Nuvaxovid COVID-19 vaccine (for ages 12 years and over)

The key safety points are:



- Approved for use for consumers aged 12 years or over receiving dose 1 and dose 2 of the primary series
- Approved for use for consumers aged 18 and over receiving the primary series and booster doses
- The Nuvaxovid COVID-19 vaccine **does not need** to be diluted
- There are 10 doses per vial
- For all vaccinator resources and materials related to Nuvaxovid please refer to the **IMAC website**.

The Nuvaxovid vaccine is available as a second line vaccine for consumers who meet the eligibility criteria.

A prescription from an authorised prescriber is required when using the Nuvaxovid vaccine as dose 2 of their primary course (i.e., following a non-Nuvaxovid COVID-19 vaccine for dose 1), in accordance with **Section 25 of The Medicines Act 1981**, as it is considered off-label use. Written consent is required for all consumers receiving an 'off label' dose of the Nuvaxovid vaccine.

## 23 Preparation of Doses

Follow the IMAC vaccine preparation instructions for vaccine preparation. These instructions are included in vaccine shipments and are available on the **IMAC website**.

**Note:** These instructions are regularly updated. Please ensure you are using the most recent version.

**Table 23.1**

Vaccine type	Dilution required?	Draw up	Doses per vial
<b>Comirnaty 15mcg/15mcg Original / Omicron BA.5-5 (grey cap) 16+ years</b>	<b>NO</b>	0.3mL	6
<b>Comirnaty 30mcg (grey cap) 12+ years</b>	<b>NO</b>	0.3mL	6
<b>Comirnaty 10mcg (orange cap) 5-11 years</b>	YES	0.2mL	10
<b>Comirnaty 3mcg (maroon cap) 6 months-4 years</b>	YES	0.2mL	10
<b>Nuvaxovid (blue cap) 12+ years</b>	<b>NO</b>	0.5mL	10

### For vaccines that **do not** require dilution:

Incorrect volume of vaccine may be detected by identifying you have drawn up less or more than **6 doses (Comirnaty)** OR **10 doses (Nuvaxovid)** from a vial.

Should this occur, immediately quarantine the vaccines and discard all doses from that vial if it is clear why the mistake has occurred.

If it is unclear why the error has occurred, keep the vaccines in quarantine and contact IMAC for clinical guidance. This error must be documented as waste in CIR and reported as an incident in the local organisation's quality and safety reporting system.

### For vaccines that require dilution (3 mcg and 10 mcg only):

Pfizer-BioNTech COVID-19 vaccine should be brought to room temperature prior to dilution, as noted in IMAC's preparing vaccine instructions. It should not feel cold to the

touch. The actual time to get the vial to room temperature will vary depending on when you take vials out of the fridge and the temperature of the room. Approximately 30 minutes should be sufficient time.

The Medsafe data sheets confirms ten (10) doses per vial.

To avoid the Comirnaty vaccine being under or over diluted it is recommended that all doses are drawn up into syringes following dilution and double-checked by a second appropriately trained vaccinator.

If the number of doses drawn from the vial are not in line with expected number this will immediately alert to the vial having not been correctly diluted. Any vial where doses drawn up are less than or more than ten (10), should be quarantined. Use of the IMAC dilution record spreadsheet will also provide an additional check.

Please discuss with IMAC on 0800 466 863 and if advised to discard, this must be documented as waste in CIR as per guidelines and reported as an incident in the local organisation's quality and safety reporting system.

## Before preparation of vaccine check:

- **That it's the right vaccine**
- Manufacturer's vaccine expiry date
- Appropriate supplies are used. **Please refer to Section 9: Vaccine and consumables ordering and delivery** for ordering consumables.

Number the vaccine vial and enter the number into the **dilution record**. Second person independently checks that the correct vaccine has been selected, by confirming the product name on the vial and checking the expiry date printed on vial by manufacturer. Second person also independently checks the numbering of the vial and documents these checks by signing/initialling the **dilution record**.

Syringe labels are used to help differentiate between vaccines. Please see **table 8.2** for the different syringe labels available to order.

During the preparation of the vaccine both expiry dates must be double checked. This includes the vial and the 10-week removal from ULT expiry date (in-use' expiry date label on the vaccine pack). Vaccines can be administered until the end of the expiry day.

**For quality and safety purposes, it is recommended that each vial and/or syringes (made from that vial), are labelled with the:**

- date and time
- expiry time

**For vaccines that require dilution (3 mcg and 10 mcg only):**

- diluent name
- date and time of dilution
- expiry time after dilution

## Vaccine preparation precautions



- Draw up from one vial at a time. Each vaccine dose from that vial should go into one kidney dish/ container with the empty vial for vaccine administration.
- It is recommended that a suitable covering is used when storing the drawn-up syringes. This is to ensure that **the vaccine is not exposed to direct sunlight or UV light** (both in the vial or in the drawn-up syringe) and that used syringes will not be put back with the unused syringes.
- During the preparation of the vaccine standard local IPC policies should be followed.
- **Any vaccine not used within the expiry time outlined above must be discarded.**
- The vaccine must not be shaken during preparation.
- Some liquid may remain in the vial after withdrawing the final dose. The leftover vaccine must be discarded. **Do not mix doses from different vials.**
- If during the preparation of the vaccine a foreign body (such as a black particle) or discolouration is identified, the vial should be discarded and recorded as an open vial-quality issue in CIR. The vaccine will appear colourless to slightly yellow, clear to mildly opalescent.
- **Note:** Call IMAC for clinical advice if required at any stage of preparation.

# 24 COVID-19 vaccine pathway to vaccination

For more information see **IMAC guidelines** found on the IMAC website and the **Immunisation handbook Section 2.2** for the correct vaccine administration process.


Please refer to the '7 Rights of COVID-19 Vaccine Administration' on the **IMAC website**.

**Table 24.1 – pre-vaccination greeting and verify identity**


Step	Action
 <p><b>Greet consumer, conduct COVID-19 health check</b></p>	<p>On arrival at the vaccination site, the vaccinator/site administrator will greet the consumer and ask whether they have any COVID-19 symptoms as per standard site practices.</p> <p>If the consumer is underage, a parent, legal guardian, caregiver, or person with an enduring power of attorney will need to accompany a child to their appointment(s) as the responsible adult and be able to provide consent for them to be immunised.</p> <p><b>Please note:</b></p> <ul style="list-style-type: none"> <li>• People who have a confirmed COVID-19 infection, should not be vaccinated until they have had the appropriate recovery period</li> <li>• People who have symptoms of COVID-19 should be advised to stay at home and get a test. They can be vaccinated once they have a negative test result and symptoms are mild only.</li> <li>• People who live with someone who has COVID-19 are a household contact and are advised to follow the specific advice public health advice for testing and isolating.</li> <li>• People who are significantly unwell are advised to wait until they are better before getting the vaccine; however, note that mild symptoms are not a contraindication. People in this situation are advised to discuss their symptoms with their GP or vaccine provider.</li> <li>• People who have been advised to self-isolate, stay at home, are under an isolation order or are waiting on a test result, should have their appointment deferred.</li> <li>• Please see the <b>Vaccination Site screening questions</b> below for questions related to clinical assessment.</li> </ul>
 <p><b>Verify consumer's identity</b></p>	<p>The vaccinator/site administrator will also verify the consumer's identity using name, DOB, address, and locate their record in CIR. This should be done in a private and confidential manner and should not be overheard or viewed by other consumers.</p> <p>Check the consumer's DOB and confirm age and what vaccine they will be receiving. If underage do <b>not</b> vaccinate.</p> <ul style="list-style-type: none"> <li>• Check with the consumer and CIR to ensure they are eligible for their vaccine today.</li> </ul>

	<ul style="list-style-type: none"> <li>• Check the dose interval and timing is correct for the vaccine the consumer is receiving. For more information see the <b>COVID-19 immunisations policy statement</b>.</li> </ul> <p><b>Note:</b> Photo ID is <b>not</b> required to confirm the consumer’s identity.</p> <p>Use the <b>7 rights of covid vaccination</b> resource available on the IMAC website.</p>
--	---

**Table 24.2 – pre-vaccination provide collateral**

Step	Action
 <p>Lead: Vaccinator</p> <p><b>Provide collateral</b></p>	<p>The vaccinator/site administrator will provide the consumer with the COVID-19 vaccination information and consent pack, which includes:</p> <ul style="list-style-type: none"> <li>• What you need to know about the COVID-19 vaccination</li> <li>• After the COVID-19 vaccination</li> </ul> <p>Ensure the consumer retains this information in either paper form or by taking a photo.</p> <p>You may also choose to provide the COVID vaccine FAQs sheet, which is available on <b>the Ministry’s website</b>.</p> <p>You may also display the privacy statement in the reception area as well as supplying the information in hard copy.</p>


**Table 21.3 – vaccination process: pre-vaccination clinical assessment**


Step	Action
 <p>Lead: Vaccinator</p> <p><b>Complete a pre-vaccination clinical assessment</b></p>	<p><b>Pre-vaccination clinical assessment</b></p> <p>The vaccinator undertakes a pre-vaccination clinical assessment. This encompasses whether the consumer has medical reasons why they should not receive the vaccine, any history of allergy, whether they had an adverse event after receiving a previous dose of the COVID-19 vaccine, any current symptoms, are pregnant or breastfeeding, and other relevant precautions.</p> <p>This includes checking that the consumer is not underage for the vaccine they will be receiving, and they have scheduled the correct interval between doses.</p> <p>For more information on dose intervals and timing see the <b>COVID-19 immunisation policy statement</b>.</p> <p><b>Interaction with other vaccines</b></p> <p>If possible, the COVID-19 vaccination should be given 7-days before or after administering the live-attenuated shingles vaccine (Zostavax). Other vaccines on the National Immunisation Schedule can be given before, after or at the same time as the COVID-19 vaccination.</p>





Step	Action
	<p><b>Boosters</b></p> <p>If the consumer has presented for a COVID-19 vaccine booster, they must meet the <b>eligibility criteria</b> available on the Ministry of Health website.</p> <p>The outcome of this clinical assessment must be recorded in CIR (in the medical screening section).</p> <ul style="list-style-type: none"> <li>• If recording the consumer as medically unfit to receive the vaccine, CIR will prompt to either cancel or reschedule the immunisation event. If the consumer is temporarily unable to receive the vaccine (that is, they are unwell today), select reschedule to ensure you can use the same CIR case record in future to capture details of the first and second doses.</li> <li>• Only select cancel if the consumer will <i>never</i> be able to receive the vaccine. Cancelling the event record means it will not be possible to go back to record a first or second dose on this record in future.</li> </ul>


**Table 24.4 – vaccination process: informed consent**

Step	Action
 <p>Lead: Vaccinator</p> <p><b>Obtain informed consent</b></p>	<p><b>Obtain informed consent before administering the vaccine</b></p> <p>The vaccinator (or vaccinator support person) <b>must</b> obtain the consumer’s informed consent to receive the vaccine prior to the administering of the vaccine.</p> <p>Where appropriate, consent may be given by a proxy such as a guardian or person with power of attorney.</p> <p>A parent, legal guardian, caregiver, or person with an enduring power of attorney will need to accompany a child to their appointment(s) as the responsible adult and be able to provide consent for them to be immunised.</p> <ul style="list-style-type: none"> <li>• If a child presents to their vaccination with whānau who cannot provide consent for the child to be immunised, written or verbal consent should be obtained from a parent, legal guardian, or person with an enduring power of attorney prior to administration of the paediatric vaccine.</li> </ul> <p>If off-label use of the vaccine, obtain <b>written</b> informed consent before administering the vaccine.</p> <p><b>Note:</b> IPC guidance must be observed when dealing with hard-copy consent forms and obtaining consent. For example, consumers should use hand-sanitiser before or after handling a pen to sign the form or bring along their own pen.</p>


Step	Action
 <p>Lead: Vaccinator</p> <p><b>Record consent in CIR</b></p>	<p><b>Consumer consent record</b></p> <p>The vaccinator or an administrative support person must record the consumer's consent to receive the vaccine in CIR.</p> <ul style="list-style-type: none"> <li>• Do not vaccinate if the interval is less than <b>21 days</b>.</li> <li>• If the person does not wish to receive the vaccine, record their decline in CIR.</li> <li>• Do not vaccinate with Nuvaxovid if the child is under the age of 12 years.</li> <li>• Do not vaccinate with Nuvaxovid if the consumer is pregnant.</li> </ul>


**Table 24.5 – vaccination process: administering the vaccination**

Step	Action
 <p>Check Vaccine</p> <p><b>Check the vaccine</b></p>	<p><b>Check the vaccine</b></p> <p>Check:</p> <ul style="list-style-type: none"> <li>• The label and confirm that you have the correct vaccine, and that the vaccine has not expired.</li> <li>• The opened/punctured diluted vial is used within the appropriate time frame before expiry. Refer to the <b>IMAC vaccine preparation sheets</b> for vial expiry times after opening.</li> <li>• The unopened vial fridge expiry date (in-use' expiry date label on the vaccine pack).</li> </ul>
 <p>Lead: Vaccinator</p> <p><b>Administer vaccination</b></p>	<p><b>Administer the vaccination</b></p> <p>Before administering the vaccine verbally check the vaccine type with the consumer. Please refer to the '7 Rights of COVID-19 Vaccine Administration' on the <b>IMAC website</b>.</p> <p>When administering concomitant vaccines, the vaccinator should ensure that the vaccines do not require any spacing and there is no specific information required to be given to consumers regarding this.</p> <p><b>Note:</b> Vaccinators should ensure the correct needle length is used for the administering the vaccine based on individual consumers being vaccinated. This includes considering body size and site vaccine will be administered (eg deltoid or vastus lateralis).</p> <p>For more information on needle length, refer to the <b>Immunisation Handbook</b>.</p>

Step	Action
 <p>Lead: Vaccinator</p> <p><b>Record information</b></p>	<p><b>Record vaccination information in CIR</b></p> <p>Once the vaccination is complete the vaccinator or administrative support person must update the consumer’s record in CIR with complete and accurate record of the vaccination event.</p> <p>This enables accurate data for operational reports (such as number of vaccinations completed and other trend data).</p> <p>This must include:</p> <ul style="list-style-type: none"> <li>• The batch, sub-batch number and expiry date for the vaccine (for example AB1234-567 the first part is the batch number, the second part is the sub-batch number) these are found on the vaccine pack.</li> <li>• The batch number and expiry date for the diluent (these are found on the diluent vial/ampoule).</li> <li>• Details of the injection site and the date and time of the vaccination event.</li> </ul> <p>In situations where this is not possible, such as CIR being unavailable, or insufficient internet connectivity at the vaccinating location, ensure an administrative process is in place to enter information into CIR on the same day as the vaccination event. This is essential clinical information; it is a requirement to ensure it is not lost and that it is transcribed correctly.</p>

**Table 24.6 – vaccination process: after vaccination**

Step	Action
 <p>Lead: Consumer</p> <p><b>Consumer waits 15 minutes in observation area</b></p>	<p><b>Observation</b></p> <p>The consumer must remain on site under observation for at least 15 minutes. If the vaccinator determines it necessary, they may ask the consumer to wait for longer than 15 minutes, for example, if the individual is in a rural or remote area or has a history of anaphylaxis. If a consumer is required to wait 30 minutes the vaccinator should record this on the CIR so that the staff member observing is aware.</p> <p>Consumers may also be required to stay for a longer length of time (20 mins) if a non-COVID-19 vaccine is to be administered at the same time eg a childhood vaccine such as MMR, shingles or tetanus booster,</p> <p>Post-vaccination advice should be given to consumers both verbally and in writing. Site Clinical Leads should ensure the latest leaflets are being used (these can be downloaded from the drop box). More information and resources can be found on the Ministry’s ‘COVID-19 vaccine: After your vaccination’ poster found on <b>the Ministry’s website</b>.</p> <p>For further information on post vaccination, see <b>section 2.3</b> in the <i>Immunisation Handbook</i>.</p>

Step	Action
 <p data-bbox="260 501 488 533"><b>Record exit in CIR</b></p>	<p data-bbox="571 295 927 327"><b>Consumer exit time record</b></p> <p data-bbox="571 340 1318 546">The site administrator/vaccinator must record the time of the consumer's exit from the site in CIR. If the consumer insists on leaving early the Site Clinical Lead must be notified and discuss this with the consumer to understand possible implications. A note should be added to the CIR to document the consumer leaving early and the advice given.</p> <p data-bbox="571 602 1286 703">Any hard copy forms must be entered into CIR by close of business on the following day. Ensure any printed copies are locked away when not in use.</p>

## 24.1 Sharing information on the vaccine

The Medicines Regulations 1984 requires written information is provided in the form of a data sheet, available at <https://www.medsafe.govt.nz/medicines/infosearch.asp>; the COVID-19 vaccine data sheet can be found by searching 'COVID-19'. There is no legal requirement for any hard copy data sheets or medicine packaging inserts to be provided on site.

## 24.2 Observation following vaccination

Consumers should remain under observation for at least 15 minutes following vaccination in an observation area. This is to ensure that any adverse reactions that may occur can receive prompt treatment.

Consumers may also be required to stay for a longer length of time (20 mins) if a non-COVID-19 vaccine is to be administered at the same time eg a childhood vaccine, shingles or tetanus booster,

All vaccinators must be able to distinguish anaphylaxis from fainting, anxiety, immunisation stress-related responses, and breath-holding spells and seizures. For further information on post-vaccination procedures, see **section 2.3** in the *Immunisation Handbook*.

### Active monitoring: Post Vaccine Symptom Check

As part of the pharmacovigilance activities for the Comirnaty vaccine, the Programme conducts active monitoring for side effects after vaccination. This is called Post Vaccine Symptom Check (PVSC) and is an SMS text-based survey sent to a randomly selected sample of the vaccinated population. When a PVSC campaign is active, a message enquiring if the consumer has experienced side effects since the vaccination is sent to up

to 25% of consumers (or their caregivers). The consumer can reply YES or NO – or reply STOP should they wish to opt out of the survey. Where the consumer replies with YES, they are sent a unique and secure link to a mobile-friendly survey form that captures the side effect/s experienced. Results are published on the Medsafe website.

There are no COVID-19 PVSC campaigns currently active.

## 24.3 Consumers' record of vaccination

Consumers should be supplied with a COVID-19 Vaccination record card detailing the vaccine administered and the date their second dose is due. This card is not designed as a vaccination certificate – and as such, may not be recognised as proof of vaccination by other countries.

### International Travel Vaccination Certificate

Consumers can request an International Travel Vaccination Certificate required when travelling overseas. This certificate can also be requested through **My Covid Record** or calling 0800 222 478.

For more information please see the **Ministry's website**.

# Section C: Additional Programme guidance, variations and incidents

Section C: Additional Programme guidance, variations and incidents - summary of changes

Version	Date	Section	Summary of Changes
51.0	28/02/23		No changes

## Section guidance

This section provides additional guidance to vaccination, NIBS, and incidents.

## Purpose

It is designed to provide additional Programme information and support, to help maintain public safety and ensure consistent and equitable vaccination outcomes across New Zealand/Aotearoa.

### Appendices relevant to this section

- **Appendix F: Links to NIBS**

# 25 Vaccination in high-risk or screened 'positive' consumers

The following is operational guidance for vaccinating consumers who are considered high-risk for being exposed to COVID-19 and are willing to be vaccinated.

While this is not advised as a general delivery model to unknown consumers, in the context of community transmission, it is important to have guidance to support this service.

'Screen positive' means that they have answered yes to any of the standard COVID-19 risk assessment/screening questions asked at vaccination reception (see Appendix G).

**Note:** There is an exception to this. Consumers with confirmed or probable COVID-19 infection **are not** recommended to be vaccinated. This reflects the lack of benefit of vaccination in this circumstance, and also risk of transmission. There is advice in the Immunisation Handbook or through IMAC to guide timing for subsequent vaccination in this scenario.

Consumers considered high risk for being exposed to COVID-19 are not suitable to be vaccinated according to the usual service design model (physical set-up of vaccination sites, workforce, and PPE guidance) as these settings are designed to be a low-risk environment. Vaccination of screen positive consumers requires additional considerations (as outlined below) as is currently recommended in only a home visit context, or in a controlled healthcare facility.

**Note:** Using this type of consumer screening, is to ensure a safe vaccination process of vaccination sites or events.

It is recommended that this section should be used in conjunction with:

- The Immunisation Advisory Centre's
- Ministry of Health **National Standards for Vaccine Storage and Transportation for Immunisation Providers 2017**.
- 2021 Addendum to **National Standards for Vaccine Storage and Transportation for Immunisation Providers 2017**.
- National Immunisation Programme Operating Guidelines.
- Ministry of Health's Immunisation Handbook 2020.

Local Health District Standard Operating Procedures. There are three scenarios below that providers could consider for the 'vaccination in high risk/screen positive consumers'.

Additional scenarios could be utilised as long as the appropriate IPC considerations are made. See additional information found “COVID-19: Infection Prevention and Control Recommendations for Health and Disability care workers”

## Scenario 1: Home Vaccination

In addition to above it is recommended that providers have Standard Operating Procedures (SOP) specific for home vaccination to support safe delivery processes.

Home visits for vaccination may be required for consumers who are unable to leave their residence because they have been required to isolate (i.e., attendance at a location of interest or contact of a confirmed case). It may also be required for those who have barriers to access due to mobility, disability, comorbidity, or another reason that means they are unable to access vaccination at a site including improving equity.

Outside the scope of this section are additional considerations which would likely be part of a Health District standard operating procedure (SOP). This could include but is not limited to a SOP on vaccine transportation and administration, staff requirements and medical emergency equipment.

## Scenario 2: Controlled Healthcare Facility

Vaccination for screen positive consumers and/or accompanying whānau in a controlled healthcare facility may be appropriate. This should only be performed in a controlled healthcare facility, where the flow of consumers and staff is controlled, such as a Hospital Emergency Department or General Practice Clinic.

This excludes dedicated vaccination sites and other settings where there is a risk of uncontrolled flow of people and workforce who are not in appropriate PPE, and so therefore is a transmission risk with other consumers and staff.

## Scenario 3: Drive-Through Vaccination

Vaccination for screen positive consumers and/or accompanying whānau in a drive-through vaccination centre may be appropriate. This should only be performed in a planned outdoor site where the flow of cars, consumers and staff is controlled. Post vaccination it is recommended they stay in their car away from others.

This excludes settings where there is a risk of uncontrolled flow of people and workforce who are not in appropriate PPE, and so therefore is a transmission risk with other consumers and staff.

PPE requirements would be the vaccinator, staff, consumer, and others in the car to wear a medical mask.



## Requirements for Scenario 1 & 2

In addition to usual vaccination processes, the following table is the requirements for the scenarios above.

	Screen Positive Requirements for Scenario 1 & 2
<b>Location</b>	<ul style="list-style-type: none"> <li>Only pre-arranged home vaccination or vaccination in a controlled residence or healthcare facility.</li> </ul>
<b>Workforce</b>	<ul style="list-style-type: none"> <li>Staff must be fully immunised.</li> <li>Home visit must have at least one authorised vaccinator and one staff on site has a CPR certificate and adrenaline administration certified. Limit staff in enclosed environment where practical</li> </ul>
<b>PPE</b>	<ul style="list-style-type: none"> <li>Consumer: Must wear a medical mask (these could be provided). All staff: P2/N95, eye protection, gown, and gloves.</li> </ul> <p>*In 'screen positive' environments, where there may also be 'screen negative' consumers, e.g., during a home vaccination, all consumers in this environment should be treated as 'screen positive'.</p> <p>**In home environments, staff should change PPE if they are moving between different houses.</p> <p>***Donning and doffing PPE outside in a home environment requires an appropriate space and transporting contaminated PPE back to base for proper disposal, this may be covered in the Health District SOP.</p>
<b>Physical Environment</b>	<ul style="list-style-type: none"> <li>Review the physical environment and consider ventilation is adequate. Discuss with local Health District IPC team if unsure.</li> </ul> <p><b>Home vaccinations</b></p> <ul style="list-style-type: none"> <li>Vaccination outside the home wherever practically possible and weather permitting. This could include in a carport, open deck area, or in their parked car. Ensure they can be observed appropriately.</li> <li>If the environment/location does not have mechanical ventilation, improve ventilation through dilution (i.e., opening windows and doors to outside air).</li> <li>If completing vaccination indoors, use a room with at least one window and keep the window(s) open for as much time as possible (outdoor temperature and safety permitting).</li> </ul> <p><b>Healthcare Facilities</b></p> <ul style="list-style-type: none"> <li>Please see section '<b>Environmental considerations and safety controls at the vaccination site</b>'. Adequate ventilation (mechanical, natural or hybrid) of all areas, including the screening, waiting, post-vaccination observation, and vaccination areas. Where a mechanical ventilation system is operating in these areas, the ventilation rate should be six air changes per hour or according to national or local requirements for healthcare facilities.</li> <li>Some older facilities may not meet the ASHRAE Standard. It is then recommended they discuss ways to improve ventilation with their local Health District IPC team.</li> </ul>

## 26 Third primary dose for severely immunocompromised

A third primary dose is recommended for severely immunocompromised consumers. It is evident that some severely immunocompromised people do not mount a sufficient immune response to provide adequate protection against COVID-19.

Advice for clinicians on the guidance is available through the Immunisation Advisory Centre, and this information will be updated periodically through the Immunisation Handbook. Clinical judgement should be applied by the prescriber to determine whether a third primary dose is required for conditions or medicines that are not listed that are associated with severe immunocompromise.

For information on the requirements for eligibility and timing, refer to **the COVID-19 immunisation policy statement**.

**Note:** There is information available on the Health Pathways site under COVID-19 Vaccination > Supporting the decision > Medical Conditions > Immunocompromised.

## 27 Vaccination and Surveillance Testing

The following section is operational guidance for providers who may wish to perform surveillance testing and vaccination at the same site, for the same consumer.

While this is not advised as a general delivery model, it is important to have guidance to support this service in the context of widespread community transmission.

Surveillance testing for COVID-19 has been used to identify cases in a community where there may be a concern around undetected transmission and infection. This would be particularly relevant in the context of a small 'community of risk' where there may be a need to both test and vaccinate consumers within a short timeframe and with an overlapping workforce.

There are differences between the processes of vaccination and testing, even in low-risk groups. Swabbing for COVID-19 is a higher transmission procedure (potentially droplet producing) than vaccinating and thus has additional PPE requirements and recommendations around physical distancing, as well as encompassing the process for swab labelling and sending to a lab.

In addition to any operational guidance, it is recommended that providers have Standard Operating Procedures (SOP) specific for vaccination and surveillance testing to support safe processes.

Due to the complexity of this process, **this model requires approval and support via NPHS Te Whatu Ora's Clinical Quality and Safety team.**

### [Requesting approval to set up](#)

Contact the NIP regional account manager to request approval to set up a vaccination and surveillance testing model.

# 28 Vaccination in Hospital

## 28.1 Introduction

The following is guidance for vaccinating consumers (including whānau of patients) against COVID-19 in a hospital setting.

Vaccination in hospital offers an opportunity to reach those who may not otherwise have access to vaccination.

Providing this service should be in accordance with local standard operating procedures, and consider local logistic, dispensing, and clinical requirements.

Consumers and/or whānau are not required to stay in hospital for the purpose of vaccination.

## 28.2 Screening

Screening for COVID-19 follows the same process outlined elsewhere in the Operating Guidelines, however the location and timing would need to be in accordance with local guidance.

Consumers that are 'screen negative' means that they have answered 'no' to all the standard COVID-19 risk assessment/screening questions (see Operating Guidance Appendix G). This means that the consumer is considered low risk for being exposed to COVID-19 and providers can follow the standard vaccination process outlined elsewhere in the Operating Guidelines.

Consumers that are 'screen positive' means that they have answered yes to any of the standard COVID-19 risk assessment/screening questions (see Operating Guidance Appendix G). This means that the consumer is considered high risk for being exposed to COVID-19 and providers should follow the Operational Guidance section "Vaccination in high-risk / screened 'positive' consumers".

## 29 Mobile vaccination team

### 29.1 Setting up mobile vaccination teams

You may choose to deliver vaccinations using a mobile vaccination team who will attend a number of different locations rather than being based at a single site. For example, this may be how you deliver vaccinations in aged residential care settings or workplaces.

As for fixed vaccination sites, you will need to consider how many vaccinators and administrators are needed for each mobile vaccination team.

### 29.2 Setting up in CIR

Mobile vaccination teams must be correctly set up in CIR, so they are linked to a facility and to track the vaccinations the mobile team have delivered.

#### **Steps to set up mobile vaccination teams in CIR:**

Complete the COVID-19 facility and site set-up details form (the regional account manager can provide a copy), with the following information:

- List each mobile team separately using a standard naming convention to identify these as mobile teams and to enable NPHS Te Whatu Ora to identify the Health District or provider linked to the team. For example, use a naming convention such as: Canterbury Health District outreach 1, MedPro mobile 2, or Auckland Health District team 3.
- Identify the facility/facilities that will be the parent for the mobile team/s.

Send the completed form to the regional account manager.

NPHS Te Whatu Ora will load the facilities and sites into CIR so users can select them. Ensure each mobile team member required to access CIR knows which facility/site they belong to. When users create vaccination events in CIR, they'll need to ensure each event record is correctly linked by checking the related contacts field under site/facility.

# 30 Home vaccinations

Vaccines can be delivered in or near a consumer's home or place of residence when they are unable to attend a vaccination site.

When administering a vaccine in a consumer's home, providers must meet the minimum requirements to safely administer the vaccine. This includes meeting the **National Standards for Vaccine Storage and Transportation for Immunisation Providers 2017** and the *COVID-19 Vaccine and Immunisation Programme Service Standards* throughout the entire process.

Providers must have a home vaccination delivery plan that includes standard operating procedures (SOPs). Prior to home based vaccinations being implemented, the plan must be approved by the Health District's immunisation clinical leads and the associated lead professional advisors.

## 30.1 Transportation of vaccine for household vaccinations

Due to regulatory restrictions on compounding and manufacturing of medicines (see section '**Transportation of diluted or drawn-up vaccine**', if a provider is utilising home vaccinations usually only one vial of vaccine can be transported and administered on each trip. This means that for each trip, the vaccinator can only transport the minimum number of doses required to vaccinate the household. This is an important consideration when planning for home vaccinations **Medicines Act 1981**. This restriction on number of vials/doses does not apply to mobile vaccination services as these will have the required resources on board to support dilution and draw up on site see section '**Mobile vaccination team**' above. All transportation of vaccine regardless of whether it is diluted or not should meet the **National Standards for Vaccine Storage and Transportation for Immunisation Providers 2017**.

The home vaccination or mobile delivery plan and SOP must cover the following:

- Maintaining staff and consumer safety, privacy, and well-being
- Respect to the consumers home and whānau
- Processes to mitigate the risk of cold-chain breaches
- Safe vaccine preparation and administration. It is recommended that preparation is carried out back at an approved vaccine preparation site. However, if not possible, preparation in a person's home should follow correct processes (i.e., double checking processes).
- Process to minimise waste
- Documentation and use of CIR
- Management of AEFI in a home environment including the immediate availability of adrenaline and phone access to call emergency services

- Operations at raised alert levels
- Risk register associated with home vaccine delivery

## 30.2 Consumer Considerations

The preferred method of vaccine delivery is at a fixed COVID-19 vaccination site. Providers should have a process to appropriately identify and approve consumers for vaccine delivery in their home.

Considerations should include:

- Consumer normally has their medical care provided in their home or place of residence.
- Does not normally leave their home or place of residence.
- Not able to be safely transported from their home to a vaccination site.
- Transport to vaccination site requires significant logistical requirement, such as multiple staff and equipment to aid transfer.
- Consumer would benefit from a home vaccination due to a disability barrier to receiving a vaccination at a site.

# 31 COVID-19 Trial Vaccinations

The Programme recognises the importance of representing the Aotearoa New Zealand population in international clinical trials and values their contribution to new COVID-19 vaccine research. **The COVID-19 Trial Vaccinations and Vaccination Certificates Policy Statement** provides a policy statement on the decisions and implications for consumers considering or participating in Medsafe approved COVID-19 clinical trial in Aotearoa New Zealand.

# 32 National Immunisation Booking System

## 32.1 Introduction

The National Immunisation Booking System known as **Book My Vaccine** supports a national-led approach to immunising New Zealand/Aotearoa against COVID-19. **Book My Vaccine** supports vaccination sites down to Community Hub level. Use by primary care sites is optional where they only service their own enrolled populations.

This section provides an operating guide for **Book My Vaccine**, including the key stakeholders, staff roles, systems, processes, and guides related to running the **Book My Vaccine** tool.

This section should be used as the first point of reference for all **Book My Vaccine** related activities by any staff member responsible for running vaccination sites and managing bookings. A detailed guide, including process flows, is available in the Detailed Booking System Guidelines document). Links to this document, training and user guides are provided in **Appendix F**.

## 32.2 Booking system principles

The **Book My Vaccine** operating model is based on the four guiding principles shown below, regarding responsibility and Governance between NPHS Te Whatu Ora, Whakarongorau Aotearoa (Whakarongorau), Health Districts and providers. These principles are intended to promote consumer safety, equity, and trust in the system. These are detailed in the four steps below:

### 1 Setup

- The **Book My Vaccine** tool supports the nationally led and locally delivered vaccination Programme.
- NPHS Te Whatu Ora has overall coordination and monitoring responsibility, including key messaging, and leading nationwide booking campaigns.
- Health Districts and providers are responsible for vaccinating their populations, including localising their campaigns to meet vaccination targets.

### 2 Setup

- The **Book My Vaccine** tool will be implemented by all Health Districts.
- The **Book My Vaccine** tool will be the trusted source of available booking slots for the public, the Health Districts and for Whakarongorau call centre to see what appointments are available for booking.
- All vaccination site types down to Community Hub level will use the **Book My Vaccine** tool. General Practices who only service their own enrolled populations will have the



option of using either their own system for vaccination scheduling or the **Book My Vaccine** tool. General Practices who service customers in addition to their own enrolled populations should use the **Book My Vaccine** tool. Pharmacies may either use their own booking system or the **Book My Vaccine** tool.

### 3 Pre-event

- The **Book My Vaccine** tool will be provided as a package with Whakarongorau as the National Call Centre
- Whakarongorau will only support the **Book My Vaccine** tool and no other booking systems once the **Book My Vaccine** tool is operational. Legacy booking systems will be phased out or replaced.
- Whakarongorau will provide a consumer supporting role for public queries (inbound) and assisted booking for all Health Districts and sites available on the **Book My Vaccine** tool.
- NPHS Te Whatu Ora is responsible to analysing booking system failures (failsafe) and developing operational process, guidelines including communication with all stakeholders.

### 4 Post-event

- The management of following up individuals for missed vaccination appointments will be a mixed model.
- Whakarongorau can provide the follow-up service for missed appointments (outbound calling) if agreed with the Health District before passing on to the Health District teams for intensive outreach follow-up. This agreement will be defined between the Health District and Whakarongorau in the engagement plan.
- Otherwise, Health District will follow-up on missed appointments (outbound calling), or they can be supported by local models with PHOs or Iwi providers.

## 32.3 Book my Vaccine system roles

The following key roles have been identified to support the **Book My Vaccine** tool. These roles include staff from the vaccination site, Health District, NPHS Te Whatu Ora and Whakarongorau. Further information related to the expected support, behaviour and outcomes of these roles is detailed above in the **roles and responsibilities table**.


**Table 22.1 – Book my vaccine tool key roles**

Key roles	Role description
<b>Site receptionist</b>	The site receptionist manages on-site check-in procedures and performs health checks prior to vaccination. The site receptionist is provisioned the role of concierge in the <b>Book My Vaccine</b> tool.
<b>Whakarongorau Aotearoa national call centre advisor</b>	The Whakarongorau national call centre advisor is the inbound point of entry for all booking queries. They are responsible for assisting consumers with creating, cancelling, and amending bookings, completing follow-up activities where commissioned to do so, and answering general vaccine related queries. They are provisioned the role of special concierge in the <b>Book My Vaccine</b> tool.  Whakarongorau conduct outbound call campaigns based on direction from NPHS Te Whatu Ora Operations Team.

Key roles	Role description
<b>Whakarongorau Health District liaison</b>	The Whakarongorau Health District liaison is the primary point of contact for communications and escalations by Districts for booking-related processes that require Whakarongorau interactions. They will escalate issues with NPHS Te Whatu Ora operations leads and the Health District operations leads, as required. The Whakarongorau Health District liaison and details regarding how to contact them will be agreed as part of the engagement plan.
<b>Site lead</b>	The site lead manages the day-to-day operation of their site and is the primary point of communication for the site. The site lead is responsible for identifying and escalating any scheduling changes to the Health District site admin.
<b>Site admin</b>	The Health District site admin acts as the first point of escalation for managing system technical operations for the sites they oversee (one site lead per ten sites is the recommended ratio). The Health District site admin is responsible for triaging and escalating impactful (minor/major event) site schedule changes. The Health District site admin is provisioned the role of site admin in the <b>Book My Vaccine</b> tool.
<b>Health District operations lead</b>	The Health District operations lead is accountable for managing the operational activities for a Health District. Their key functions include generating reports to identify Health District follow-up activities, managing all escalations required for a Health District's attention, and sharing escalations with NPHS Te Whatu Ora and Whakarongorau as required. They are accountable for the accuracy of all site schedules.
<b>The NPHS Te Whatu Ora operations lead</b>	The NPHS Te Whatu Ora operations lead is the primary point of contact for escalations into NPHS Te Whatu Ora. Their key obligation is managing communications between NPHS Te Whatu Ora and Whakarongorau/Health Districts. They are provisioned the role of super user in the <b>Book My Vaccine</b> tool and are responsible for onboarding users and sites in the system. NPHS Te Whatu Ora operation team are responsible for failsafe reporting and organising outbound call campaigns to reach consumers.

## 32.4 Booking system processes and best practice

Figure 22.1 – booking tool processes:



## 32.5 Setup and maintenance

### Creating a new site

Creating a new site relates to setting up a site on the booking system. Health Districts are asked to nominate one individual to be responsible for sending information and communicating between parties. If possible, it is preferable that new sites are set up in batches (such as all sites going live in a two-week period are grouped together) to minimise duplication of processes.

The Health District or provider will populate the new site CIR and NIBS setup forms. The nominated Health District staff member is then required to email the forms to the NPHS Te Whatu Ora central help desk. A seven-day lead time for receiving the completed forms prior to site going live is required by NPHS Te Whatu Ora. The information within this form is used by teams within NPHS Te Whatu Ora to setup the new site in CIR and NIBS. It will take up to 72 hours for these sites to be created in the respective systems, after form submission.

New system users required for the site will complete qualifications and training processes as prescribed by their Health District workforce lead. System training includes NIBS and the Covid Immunisation Consumer Support (CICS) tool (if the Health District is using CICS, its use is optional). System users will be trained for the site admin and concierge roles in NIBS and the system user role for CICS. The nominated Health District staff member will inform NPHS Te Whatu Ora once system user training has been completed (it is mandatory for the training to be completed before users are provisioned). NPHS Te Whatu Ora will then provision the appropriate users for the site. The nominated Health District staff member will receive a confirmation of this task completion via email.

## Amend site schedule

Amending a site schedule involves updating the capacity and availability of appointment slots for a site. The site lead or site administrator is responsible for identifying major schedule changes, escalating these to the Health District operations lead, and making necessary system changes. The thresholds for escalation will be set and be maintained by each Health District.

**Note:** Changing the schedule in NIBS does not cancel or reschedule any existing bookings. Refer to the **event rebooking** section below for details.

**It is crucial the site lead or site administrator performs an impact assessment regarding bookings when amending a site schedule, specifically when the number of appointment slots are reduced.**

## Event rebooking

In the case of an event causing a disruption to a site, where an existing schedule is set, and appointments are cancelled, consumers must be rebooked in the system. Event severity, minor or major as be determined by the site and Health District staff, will dictate the applicable escalation path to ensure key stakeholders have visibility of the event, and can assist with implementing appropriate resolution measures. When a major event occurs such as a bulk cancellation or a bulk reschedule, this must be approved by NPHS Te Whatu Ora. Further information is outlined in the Detailed Booking Systems Guidelines document found in **Appendix F**.

**Note:** Rescheduling is not automatic function. Consumer appointments will not be cancelled or rescheduled when a site schedule change is created.

### Amend site details

Amending site details involves updating the site location and other site properties. These changes do not affect scheduling. The site lead, site administrator or Health District operations lead is responsible for identifying such changes are necessary. The site administrator is responsible for making the changes in the system.

## Pre-event

### Booking an appointment

Where a consumer is eligible to be vaccinated, they are able to book a primary course with the option to include two appointments at the same location, a first dose only and a second dose only (after the required number of days from their second dose). The COVID-19 vaccine is different for certain age groups and so consumers need to select the relevant age band.

When eligible, consumers can select Pfizer, paediatric Pfizer or Novavax as the vaccine type. Eligible consumers can also book a booster dose after the required number of days from when their primary course was completed and a second booster dose after the required number of days from when the first booster dose was administered.

Consumers are asked to provide their personal details, allowing the system to send a booking reference and confirmation to the consumer. A contact person's details are required for appointment confirmation and reminders where the booking is for a child aged 6 months to 11 years. Bookings can also be made by an individual on behalf of a consumer (for instance a family member or friend), or through Whakarongorau.

Consumers can select to book as a small group (2-5 people) or a large group (6-30 people) for a single dose of a Pfizer vaccine. For a small group, consumers are asked to provide their personal details and the contact details for a booking arranger who receives the booking references, confirmation, and reminders for all individuals in the group. For a large group, only the first and last name and contact information of the booking arranger are required.

**Note:** The third primary dose of Pfizer for people who are severely immunocompromised is not booked via the **Book My Vaccine** tool.

### **Update and/or cancel an appointment**

Consumers can update the time and/or location of their vaccine appointment/s or cancel their appointment/s through the **Book My Vaccine** tool. Consumers must enter either an email address or phone number. This will be used to provide the consumer with a confirmation of their booking (such as the booking reference etc.). If a consumer does not have this information, they should contact Whakarongorau for assistance. Consumers have the option to rebook or cancel an appointment any time up to two weeks after the scheduled appointment date. Consumers cannot reschedule as a small group, rather each individual should be cancelled or rescheduled using the contact details of the booking arranger and the booking reference of the individual. Large groups can only be cancelled using the booking arranger's contact details and the booking reference.

## During the vaccination event

### **Consumer arrival**

Where a consumer has booked an appointment as an individual or as part of a small group and arrives at a site, an Immunisation Case will be in CIR. The Concierge must confirm that the consumer is eligible for the Vaccination Plan and then check the consumer in. Where a consumer is part of a large group, no Immunisation Case will be in CIR and this will need to be created as per a walk-in.

Where a consumer arrives at a site without an appointment (walk-in) or if they show up early for an appointment, providing the Health District or provider has capability to take walk-in consumers and the site has availability, the consumer may be vaccinated. As no appointment was booked, an Immunisation Case will need to be created to record the vaccination activity in CIR. The Concierge must be sure to select the correct Vaccination Plan when creating a new Immunisation Case.

Walk-in consumers should be assisted to book their next appointment on BMV. This process is best practise to ensure that consumers are booked to receive the next dose.

## Post event: follow-up

### **Booking Did Not Attend (DNA) follow-up**

Health Districts remain accountable to ensure that consumers who did not attend (DNA) appointments are contacted. Health Districts may utilise Whakarongorau to undertake any follow-up activities to rebook consumers. This process to request follow-up services will be defined in the engagement plan between Whakarongorau and each Health District. This can be a mixed model, where each party may be responsible for following up with different groups of consumers within the Health District. The best practice for following-up is to contact a consumer the day after the missed appointment, followed by two more attempts, at the fourth day and seventh day. Where the consumer remains uncontactable, the Health Districts is responsible for executing the most suitable follow-up response (further outbound calls or ceasing follow-up communications).

# 33 Incidents

## 33.1 Incident management

The site team should be trained and prepared to respond to three possible medical emergencies associated with COVID-19 vaccination: fainting, hyperventilation, and anaphylaxis. The appropriate medication and equipment must be on site to manage these incidents.

Refer to **section 2.3 of the *Immunisation Handbook*** for guidance on emergency equipment required to manage post-vaccination medical emergencies.

Adverse events should be managed in accordance with HQSC ***Guide to the National Adverse Events Reporting Policy 2017***.

In the event of a serious adverse event or incident it is important to follow organisational process to report, review, and learn from the incident.

- **Appendix I** outlines the process steps for notifying serious incidents to the Programme. This includes the COVID-19 Vaccine related severity assessment codes (SAC) and the form required to notify the Programme of incident and serious adverse events.

## 33.2 Adverse events during observation period

If any consumer has an adverse event during the 15-minute observation period at the vaccination site, appropriate medical attention must be provided. The on-site adverse event must be recorded and submitted in the CIR to support reporting on adverse events following COVID-19 vaccine immunisation.

For more information regarding managing medical emergencies and anaphylaxis, please see **section 2.3 of the *Immunisation Handbook***.

## 33.3 Recording an anaphylaxis event

Where a suspected anaphylaxis event occurs following a vaccination event, it is important to record and submit consumer details of the event in the CIR. The person who handled the event must complete the anaphylaxis checklist record (found on the **IMAC website**) as soon as practical. The anaphylaxis checklist should be completed and uploaded via the Dropbox to the CARM **link**.

Adverse events should be notified to the site lead clinician, who can undertake a clinical review and determine appropriate actions with the site manager (such as pausing vaccinations for a time, should this be required).

## 33.4 Adverse events after observation period

Consumers should be advised by the vaccinator, at the time of vaccination, of common **and** rare side effects that can occur after the observation period (after they've left the vaccination site). This should include a discussion about when and how to seek medical attention, and how to submit an adverse reaction report to CARM.

**Common side effects of COVID vaccines** include pain, redness or swelling at the injection site, feeling tired or fatigued, headache, muscle or joint aches and pain, chills, fever, and nausea. These effects are usually mild or moderate and improve within a few days after the vaccination.

### **Rare side effects of COVID vaccines**

Myocarditis and pericarditis are an inflammation of the heart muscle or lining and can range from mild to serious illness. They are usually caused by viruses but are also a **rare side effect** of both the Pfizer and Novavax COVID-19 vaccines.

Symptoms of myocarditis and pericarditis linked to the vaccine generally appear within a few days, and mostly within the first week after having the vaccine. Consumers should be advised that if they get any of these new symptoms, they should seek medical help, especially if these symptoms don't go away:

- Tightness, heaviness, discomfort or pain in your chest or neck.
- Difficulty breathing or catching your breath
- Feeling faint or dizzy or light-headed
- Fluttering, racing, or pounding heart, or feeling like it is 'skipping beats'.

## 33.5 COVID-19 treatment injury claims

ACC is sharing advice with providers regarding lodging ACC claims for a physical injury resulting from a COVID-19 Vaccination. Such injuries may be covered by ACC if the injury criteria for treatment are met. Under ACC legislation, the injury must be clearly caused by the vaccination and must not be a necessary part or ordinary consequence of the treatment. For example, inflammation around the site of the injection is common with COVID-19 Vaccination (an ordinary consequence) and is unlikely to be covered. Infections (such as cellulitis or septic arthritis) due to the vaccination, and anaphylaxis resulting in injury are not ordinary consequences and are more likely to be covered.

Where a consumer has an injury that meets these criteria, they may require further treatment or support. In such cases, providers should lodge an ACC2152 treatment injury claim form with ACC as well as an electronic or manual ACC45 injury claim form. These forms and more information can be found on **ACC's website**.



Providers will need to include the vaccine brand and identifying dose number (for example, whether it the first or second Pfizer-BioNTechCOVID-19 Vaccine dose).

**Note:** Health providers should keep good clinical records of reactions and complications and arrange appropriate clinical management and follow up. Treatment injury claim forms can be completed at the time or any time after the event. However, if longer than 12 months additional information is required. Time should be taken to obtain consumer consent for a claim to be lodged with ACC, as it involves providing their personal and private information to ACC. Consumers should be reassured the health system will manage their treatment regardless of an ACC claim.

## 33.6 Recording vaccine errors

A vaccine administration error is any preventable event that may cause or lead to, inappropriate use of a vaccine or consumer harm. Administration errors can occur at any stage of the vaccination process (such as storage or handling, site/route of administration, or dosage given).

Some known vaccine errors include unauthorised age group vaccinations, shorter than recommended dosing intervals, injecting errors, dosage errors, vaccine administration errors, or when the consumer has an adverse event due to a vaccine error.

### In the event of a vaccine administration error

- Inform the consumer/s involved. This should occur within **seven working days**.
- If guidance/advice is needed, consult **IMAC on 0800 IMMUNE (466 863)**, option 1 (health professionals) and then option 2 (COVID-19 vaccinator support)
- Record the error in CIR under adverse events error to provide for reporting on vaccine administration errors.
- Determine how the error occurred to provide for strategies to be implemented to prevent a recurrence.

Providers should report all COVID-19 vaccine administration errors, including those not associated with an adverse event. Upon submitting the adverse event/medical error form to CIR, data will go to the medical assessment team at CARM. Please provide as much detail as possible about the error that occurred, any actions that were taken at the time of the event and pending actions. The medical assessment team review adverse events and medical errors to help inform any follow up required. Adverse event and medical error reports also inform vaccine safety monitoring.

## 33.7 Early second doses

If the first and second dose of the Pfizer-BioNTech COVID-19 vaccine is administered at an interval of **less than 21 days**, or **28 days** for the Nuvaxovid vaccine following another COVID-19 vaccine, this is reported as an early second dose.

**In the event of an early second dose, please follow the instructions below with respect to the reported cases:**

- Verify the case ID entry - if wrong, then correct the CIR record.
- If correct, complete a CARM medication error report as this is a 'never event' use of the vaccine.
- Inform the affected person of the error and ask them to report any reactions – refer to the handout 'After the COVID-19 vaccination'.
- Clinical advice (e.g., by the medical advisors at 0800IMMUNE) may be required. This will depend on the timing of the second dose and the characteristics of the individual.
- Identify improvements to local practice and process to avoid early second doses and share the learnings as soon as possible.
- On investigation, and if in the event the person reports possible harm, then follow your Health District or provider's adverse event process and or complaints process.
- If an adverse reaction or injury is experienced by the individual following the event, submit an additional CARM AEFI report and arrange ACC treatment injury claim per **ACC2152 form**.

# 34 Variations

## 34.1 Missing or incorrect information in the CIR

When it's identified a consumer has missing or incorrect information documented in the CIR relating to the administration of a vaccine in Aotearoa New Zealand, then it must be corrected as it is a legal record.

The CIR can be modified by the provider or health professional within 24 hours after details of a vaccination were entered.

After this time the CIR can only be modified by contacting **help@C-19imms.min.health.nz** or **0800 223 987**.

## 34.2 Where the consumer has received vaccination overseas

This advice applies when consumer has received a COVID-19 vaccine overseas (which may or may not be of the Pfizer COVID-19 vaccine):

- Specifically, when the consumer has been administered 1 vaccination of a 2-dose primary course (Pfizer or other two-dose vaccine), they are able to receive the Pfizer COVID-19 vaccine if the second dose is at least 4 weeks after their overseas vaccination.
- Please consult IMAC on 0800 IMMUNE for specific clinical advice.

The consumer must provide evidence of their overseas vaccination (e.g., a vaccine receipt card or other documentation). The provider creates a CIR immunisation record for the 2-dose vaccine and uploads the overseas first dose vaccination evidence the consumer has provided.

**Note 1:** Based on advice from the COVID-19 Vaccine Technical Advisory Group the CIR will only accept the entry of named vaccines. At the time of review of this document those overseas vaccines able to be entered in CIR are contained at: -

[www.health.govt.nz/our-work/diseases-and-conditions/covid-19-novel-coronavirus/covid-19-vaccines/my-covid-record-proof-vaccination-status/covid-19-overseas-vaccinations-and-certificates](http://www.health.govt.nz/our-work/diseases-and-conditions/covid-19-novel-coronavirus/covid-19-vaccines/my-covid-record-proof-vaccination-status/covid-19-overseas-vaccinations-and-certificates)

**Note 2:** The CIR record of the Aotearoa New Zealand-based dose administered is automatically notified to the consumer's primary care (GP) practice through the existing CIR primary care practice notification functionality. The entry into the CIR of an overseas vaccination contained in the preceding hyperlink is likewise automatically notified to the consumer's primary care practice.

# Appendices

## Appendices: summary of changes

Version	Date	Appendix	Summary of Changes
51.0	28/02/23	J	Updated table to remove outdated information

## Section guidance

This section provides the appendices for the Operating Guidelines.

### Purpose

It is designed to provide additional information and support, to help maintain public safety and ensure consistent and equitable vaccination outcomes across Aotearoa New Zealand.

# Appendix A:

## Site checklist

As a general principle, the site and staff should be prepared and adhere to standard operating policies and standards, including the clinical governance and health and safety, expected in a clinical environment to ensure staff and consumer safety.

Tables A1 to A5 below, provide an overview of the minimum requirements to deliver COVID-19 vaccinations safely and efficiently.

**Table A1 – plan checklist**

Plan	Y / N	Comments
<p><b>Vaccination volume plan</b>            Vaccination sites have planned for expected daily volumes of vaccine recipients, considering:</p> <ul style="list-style-type: none"> <li>• Staffing numbers</li> <li>• Space and distancing</li> <li>• Privacy and confidentiality</li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Workforce plan            To maintain the staff roster including managing unavailability, illness, and other absences.</p>	Y <input type="checkbox"/> N <input type="checkbox"/>	
The list of <b>Key Contacts</b> is up to date and accessible.	Y <input type="checkbox"/> N <input type="checkbox"/>	
<b>Clinical Quality and Safety</b> oversight is on site.	Y <input type="checkbox"/> N <input type="checkbox"/>	
<p><b>Local development of:</b></p> <ul style="list-style-type: none"> <li>• Infection Prevention</li> <li>• Control guidance</li> <li>• SOPs</li> <li>• Cold Chain Accreditation for this site</li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p><b>Site locations consideration:</b></p> <ul style="list-style-type: none"> <li>• Location/traffic/access/parking/signage</li> <li>• Availability of public transport</li> <li>• Accessibility (including disability access to parking and to vaccination site building)</li> <li>• Traffic management</li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
The site can maintain temperature requirements of the vaccination preparation space.	Y <input type="checkbox"/> N <input type="checkbox"/>	
A <b>documented risk assessment</b> has been conducted for every individual vaccination site including business continuity plan covering changes in COVID response alert levels	Y <input type="checkbox"/> N <input type="checkbox"/>	

Site-specific COVID Tracer App QR codes have been created.	Y <input type="checkbox"/> N <input type="checkbox"/>	
A plan is in place to maintain adequate and appropriate resources including: <ul style="list-style-type: none"> <li>• PPE supplies</li> <li>• Vaccine and consumables</li> <li>• IPC supplies</li> <li>• Waste management</li> <li>• Signage</li> </ul>	Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/>	
A plan is in place to <b>maintain daily supplies</b> of consumer collateral, including: <ul style="list-style-type: none"> <li>• What you need to know about the COVID-19 vaccination</li> <li>• COVID-19 vaccination consent form</li> <li>• After the COVID-19 vaccination</li> <li>• Privacy Statement</li> <li>• Hard-copy form to collect household contacts.</li> </ul>	Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/>	
A plan is in place for <b>equitable access</b> , including: <ul style="list-style-type: none"> <li>• Access to translation and interpretation services</li> <li>• Written material and signage in easy-to-read formats</li> <li>• Supporting resources/literature is available in a range of languages/formats for those with low health literacy.</li> <li>• Service delivery model provides for whanau/support people accompanying consumers.</li> <li>• Venue access caters for disabled people and support for those with visual or hearing impairments.</li> </ul>	Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/>	
A plan is in place to manage the transition from locally managed booking systems to the national immunisation booking system or Programme accepted booking solution.	Y <input type="checkbox"/> N <input type="checkbox"/>	
A site evacuation plan is in place.	Y <input type="checkbox"/> N <input type="checkbox"/>	
A dry run has been completed for all vaccination sites.	Y <input type="checkbox"/> N <input type="checkbox"/>	

**Table A2 – place site checklist**

Physical site	Y / N	Comments
Adequate space (including also for whanau/support persons) and associated capacity for: <ul style="list-style-type: none"> <li>• Screening</li> <li>• Registration</li> <li>• A private space for consultation, family groups, and vulnerable people requiring support</li> <li>• Waiting (seated)</li> <li>• Vaccination (including drawing up and administering)</li> </ul>	 Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/>	

<ul style="list-style-type: none"> <li>Post-vaccination observation.</li> </ul>	Y <input type="checkbox"/> N <input type="checkbox"/>	
Access to secure storage for medical records (including consent forms).	Y <input type="checkbox"/> N <input type="checkbox"/>	
Appropriate signage to identify as vaccination site for consumers, including COVID-19 vaccination campaign posters/banners/flags. Signage should also include Code of Consumer Rights.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Site has clearly marked one-way foot traffic flow with clear entry and exit areas.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Adequate number of hand-hygiene stations in strategic areas for public and staff	Y <input type="checkbox"/> N <input type="checkbox"/>	
Appropriate emergency medication, equipment, and space to respond to medical emergencies. All equipment in the site to be well maintained, in good working order, calibrated/monitored as required and with current electrical safety compliance testing/certificates as necessary. <b>Note:</b> This should also include equipment suitable for children if the site will be administering paediatric vaccines.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Appropriate cold chain provisions that are applicable for the site are in operating order, including having appropriate refrigerators and opaque containers to store supplies. Cold Chain Accreditation is held and is current if applicable.	Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N/A <input type="checkbox"/>	
Adequate space for vaccine storage and preparation.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Adequate security (e.g., alarm, overnight security guard) if vaccine is to be stored at vaccination site overnight.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Appropriate waste management facilities, including facilities in place to safely dispose of sharps and unused, damaged, or empty vaccine vials (e.g., Interwaste vial disposal bin ordered).	Y <input type="checkbox"/> N <input type="checkbox"/>	
Vaccination stations at least one metre apart.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Access to CIR-compatible IT hardware including tablets, laptops or desktop computers with screens positioned out of sight of unauthorised persons.	Y <input type="checkbox"/> N <input type="checkbox"/>	
IOS or Android smartphones with Salesforce Authenticator app available to CIR users.	Y <input type="checkbox"/> N <input type="checkbox"/>	
High-speed wireless or 4G coverage.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Access to appropriate internet browser (Note: Internet Explorer is not supported).	Y <input type="checkbox"/> N <input type="checkbox"/>	

**Table A3 – process checklist**

Process	Y / N	Comments
Scheduling of vaccination appointments avoids over-crowding and allows for physical distancing.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Booking and scheduling system includes arranging for consumers to return for a second dose of the vaccine at least three weeks after receiving the first dose.	Y <input type="checkbox"/> N <input type="checkbox"/>	
All staff have access to the Operational Guidelines.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Procedures are in place for identifying vaccine recipients.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Process in place for screening all staff for signs and symptoms of COVID-19 at the start of each shift.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Standardised screening processes are in place for contraindications, receipt of previous dose of COVID-19 vaccine or other vaccines, and COVID-19 symptoms.	Y <input type="checkbox"/> N <input type="checkbox"/>	
'Where to get help' poster is accessible to all staff.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Consumer information processes in place, including the provision of consumer collateral.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Cold chain process in place, site delivery and receipt.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Processes in place for infection prevention and control including: <ul style="list-style-type: none"> <li>• Hand hygiene</li> <li>• PPE protocols</li> <li>• Injection safety</li> <li>• Needlestick injury protocol</li> </ul>	Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/>	
Processes in place to safely manage waste and for safe disposal of sharps and unused, damaged, or empty vaccine vials.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Process in place for monitoring, managing, and reporting adverse events following immunisation, including anaphylaxis.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Policies in place for blood body and fluid exposures (BBFE) and infection prevention control (IPC).	Y <input type="checkbox"/> N <input type="checkbox"/>	
Appropriate process in place to respond to medical emergencies associated with the vaccination.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Incident management procedures are in place and staff know how to report any clinical incident.	Y <input type="checkbox"/> N <input type="checkbox"/>	
SOP available for accessing and operating CIR and completing inventory reporting requirements.	Y <input type="checkbox"/> N <input type="checkbox"/>	
Business continuity plans in place, including access to hard-copy versions of: <ul style="list-style-type: none"> <li>• Consent forms with CIR data fields on the reverse and associated secure storage.</li> <li>• COVID-19 Vaccine Adverse Event Report</li> </ul>	Y <input type="checkbox"/> N <input type="checkbox"/> Y <input type="checkbox"/> N <input type="checkbox"/>	



**Table A4 – workforce checklist**

Workforce	Y / N	Comments
<p>Staffing levels (including trained and accredited as required) are appropriate for delivering the scheduled vaccination volume. At a minimum, the following functions need to be allocated:</p> <ul style="list-style-type: none"> <li>• Consumer welcome</li> <li>• Preparation and administration of doses</li> <li>• Obtaining informed consent</li> <li>• Events recording in CIR by a CIR-trained person</li> <li>• After-immunisation observation</li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Site workforce encourages equitable access and the workforce demographic, as reasonably practicable, reflects of the likely consumer population or local area.</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Staff are educated in disability equity access and know how to apply supported decision-making approach (e.g., the Ministry's <b>Disability equity course</b>)</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Staff accessing consumer data have completed the appropriate privacy training (e.g., see the <b>Privacy Commissioner courses link</b>).</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Staff inducted to the site and to have completed all relevant training including cold chain and IMAC/vaccine training, adverse event training, and CIR training.</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Appropriate staff training to respond to three possible medical emergencies associated with the vaccination (fainting, hyperventilation, and anaphylaxis).</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Staff roles and responsibilities are clearly defined.</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Multi-vaccinator sites have a named Lead Clinician.</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>An appropriate person has been identified to receive vaccine delivery as part of cold chain provisions.</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Infection Prevention and Control staff have been identified including:</p> <ul style="list-style-type: none"> <li>• IPC Lead</li> <li>• IPC trainers</li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>Security presence available to control access to the site and be available for support in the event of attempted unauthorised access.</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	
<p>All vaccination site staff have been given the opportunity to receive a COVID-19 vaccination.</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>	

**Table A5 – other considerations checklist**

Other considerations	Y / N
<ul style="list-style-type: none"> <li>Staff working in or near MIQ or other locations that may require additional infection prevention controls, must adhere to the standard SOPs and associated protocols for such locations, including physical distancing requirements.</li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>
<ul style="list-style-type: none"> <li>In the event of a change in Alert Levels, adherence to the relevant PPE SOPs and associated protocol is required to operate under the Alert Level, including physical distancing requirements.</li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p>
<ul style="list-style-type: none"> <li>Where a mobile vaccination team is being set up, in addition to the above also consider the following:               <ul style="list-style-type: none"> <li>Staff numbers to match expected demand as well as site health and safety requirements</li> <li>Site security</li> <li>Appropriate training</li> <li>Correct set up in CIR, including completion of the 'COVID-19 Facility and Site Set-up Details'.</li> <li>Reliability of supply of resources and equipment</li> <li>Internet connectivity to enable use of CIR</li> <li>Logistics, including vaccine storage and transport</li> <li>Business continuity</li> </ul> </li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p>
<ul style="list-style-type: none"> <li>Drive through vaccinations:               <ul style="list-style-type: none"> <li>Some disabled people use modified vehicles that seat the driver/passengers higher – potentially making it more difficult for vaccinators to reach</li> <li>A reminder that car doors can also be opened if proper needle positioning can't be achieved through the window</li> </ul> </li> </ul>	<p>Y <input type="checkbox"/> N <input type="checkbox"/></p> <p>Y <input type="checkbox"/> N <input type="checkbox"/></p>

Version 3

# Appendix B:

## New facility/site setup

This information must be provided to NPHS Te Whatu Ora five days in advance of any initial deliveries. Please use the following template to complete the information required to enable us to set up a vaccination facility or vaccination site. Please take care and provide detail when completing the form, as accurate information is required to ensure successful delivery of vaccines and consumables.

Return the completed form to [help@c-19imms.min.health.nz](mailto:help@c-19imms.min.health.nz) and CC your Regional Area Manager

Version	Date	Summary of Changes
1.4	05/11/21	New facility/site setup form updated with changes to categorisation of site types and provider types.
1.5	29/03/22	Addition of flu vaccination to vaccination type field Amendment to list of 'site types' Addition of cold chain storage expiry date Addition of field to record location of back up cold-chain storage
1.6	22/07/22	New section 4 added for 'Facility move details'
1.7	23/08/22	"Vaccine type recorded in NIR" added as a further Vaccine type to accommodate inventory only vaccines that facilities can order, stock, and consume. Boostrix option added to "Vaccine type recorded in NIR field".
1.7.1	23/09/22	Pharmacy License expiry option added, as a pharmacy licence holder is deemed to hold CCA for as long as they hold a current pharmacy licence.

Previous revision history can be found at the end of the appendices section.



Has the site been signed off by the Health District CE?		Please attach a copy of signed authorisation		
Y <input type="checkbox"/>	Please tick if yes	Y <input type="checkbox"/>	Please tick to confirm	
Location details section		New site set up – part one of four		
<b>A</b>	<b>Site</b> <i>Only complete Section A if a site is being set up. Note: Sites are where vaccines are administered</i>			
<b>Site</b>	Health District	Enter the Health District in which the vaccination facility/site is located		
	Site name	Please provide the site name		
	Site address	Please provide the delivery address. Please include floor number/building number/gate number if relevant.		
	Confirm	Suburb and post code of this site		
	City	Enter city in which this site is located		
	Site type details			
	Please tick	Is this vaccination site also a facility? Y <input type="checkbox"/> N <input type="checkbox"/>		
	Vaccine type (Recorded on CIR)	<input type="checkbox"/> Covid-19 <input type="checkbox"/> Influenza <input type="checkbox"/> Both Covid-19 & Influenza		
	Vaccine Type (Recorded on NIR)	<input type="checkbox"/> Boostrix <b>Please note: Use to request stock of vaccines only</b>		
	Site type Please tick	<input type="checkbox"/> GP <input type="checkbox"/> Hospital <input type="checkbox"/> Marae <input type="checkbox"/> Off-Site <input type="checkbox"/> On-Site <input type="checkbox"/> Mobile or Pop-up Site ( <i>short term vaccination site</i> ) <input type="checkbox"/> Mass Vaccination Event <input type="checkbox"/> Permanent Vaccination Centre ( <i>long term vaccination site</i> ) <input type="checkbox"/> Drive-Through <input type="checkbox"/> School <input type="checkbox"/> Community Pharmacy <input type="checkbox"/> Urgent Care Clinic <input type="checkbox"/> Residential Facilities (e.g. Aged Care Facility, Residential Care etc.) <input type="checkbox"/> Place of Worship <input type="checkbox"/> Workplace (Vaccination for staff and whanau) <input type="checkbox"/> Bus <input type="checkbox"/> Other:		
	Equity focus	<input type="checkbox"/> Not applicable <input type="checkbox"/> Māori <input type="checkbox"/> Pacific Island <input type="checkbox"/> Disability <input type="checkbox"/> Mixed		
	The following information relates to the Provider(s) responsible for the site.			
	Primary Provider name	Please provide the name of the primary provider		
	Provider type	<input type="checkbox"/> Health District <input type="checkbox"/> Occupational Health <input type="checkbox"/> Community Pharmacy <input type="checkbox"/> GP <input type="checkbox"/> PHO <input type="checkbox"/> Hauora <input type="checkbox"/> Pacific Health Provider <input type="checkbox"/> Urgent Care Facility <input type="checkbox"/> Other If other, please add details		
Provider equity focus	<input type="checkbox"/> No Specific Equity Focus (General population) <input type="checkbox"/> Māori <input type="checkbox"/> Pacific Island <input type="checkbox"/> Disability			
Collaborating provider name	Please provide the name of the collaborating provider (if applicable)			
Collaborating provider type	<input type="checkbox"/> Health District <input type="checkbox"/> Occupational Health <input type="checkbox"/> Community Pharmacy <input type="checkbox"/> GP <input type="checkbox"/> PHO <input type="checkbox"/> Hauora <input type="checkbox"/> Pacific Health Provider <input type="checkbox"/> Urgent Care Facility <input type="checkbox"/> Other If other, please add details			
Collaborating provider equity focus	<input type="checkbox"/> No Specific Equity Focus (General population) <input type="checkbox"/> Māori <input type="checkbox"/> Pacific Island <input type="checkbox"/> Disability			

<b>B</b>	<b>Facility</b> <i>Please provide Facility or Associated Facility details.</i> <b>Note: Facilities are where vaccines are shipped to, stored, and subsequently distributed to sites</b>
----------	--

<b>Facility</b>	<b>Health District</b>	Please provide the Health District where the facility is located
	<b>Facility name</b>	Please provide the facility name if different to site name in Section A
	<b>Facility type</b>	Please provide the facility type, such as hospital, pharmacy, clinic
	<b>Facility address</b>	Please include suburb, city and postcode
	<b>Delivery address (if different from facility address)</b>	Please advise the delivery address - include floor number/building number/gate number if relevant.
	<b>Facility ID (HPI ID)</b>	What is this facility's ID (if unknown, state 'unknown')

<b>Delivery information</b>
-----------------------------

Please provide the available delivery times for the facility, such as 7am to 5pm, Monday to Friday.										
<b>Available delivery times</b>	<input type="checkbox"/> Monday		<input type="checkbox"/> Tuesday		<input type="checkbox"/> Wednesday		<input type="checkbox"/> Thursday		<input type="checkbox"/> Friday	
	AM	PM	AM	PM	AM	PM	AM	PM	AM	PM
<b>Delivery Notes</b>	Please add any comments which may assist the delivery driver in successfully									

Storage, capacity, and contact details		New site set up – part three of four	
<b>C</b>	<b>Which of the following storage accreditations does the facility provide?</b>		
Ultra-cold (-70C)	Y <input type="checkbox"/> N <input type="checkbox"/>	If yes, please provide details of how many vials can be stored	
Frozen (-20C)	Y <input type="checkbox"/> N <input type="checkbox"/>	If yes, please provide details of how many vials can be stored	
Cold chain (2-8C)	Y <input type="checkbox"/> N <input type="checkbox"/>	If yes, please provide details of how many vials can be stored	
Cold chain (2-8C) accreditation or Pharmacy License expiry date	Expiry Date: [DD/MM/YYYY]		
Back-up fridge location	[Please enter name and address of alternative location]		
Ambient	Y <input type="checkbox"/> N <input type="checkbox"/>	If yes, please provide details of how many vials can be stored	
Consumables	Y <input type="checkbox"/> N <input type="checkbox"/>	If yes, please provide storage details	
Is there a data logger reader at location?	Y <input type="checkbox"/> N <input type="checkbox"/>	If yes, please provide details about brand/type	
<b>Pay per dose contract</b>			
Pay per dose contract number	If this contract is a Pay per Dose contract – Please provide the contract number.		
Regional Anniversary	In which region will you be observing Regional Anniversary days?		
<b>Pay per dose contract</b>			
Named role	Please confirm the <b>named role</b> at this vaccination facility/site who will be available and is authorised to receive the vaccine/consumables upon delivery, for example lead nurse, clinic manager.		
Named role Name and contact phone number/s	Name	Confirm name	
	Phone	Confirm phone number/s	
Alternate Name and contact phone number/s of other team members who fit the named role	Name	Confirm name alternate 1	
	Phone	Confirm phone number/s alternate 1	
	Name	Confirm name alternate 2	
	Phone	Confirm phone number/s alternate 2	
<b>Completed/signed by</b>			
Name	Add name		
Title	Add title		
Signature	Insert signature		

Location details section		Facility moving – part four of four
<b>D</b>	<b>Only complete Section D if a facility is moving (e.g. due to an expired lease), <u>and</u> Cold Chain Accreditation can be transferred. If Cold Chain Accreditation cannot be transferred, please complete a site closure and new site set-up form.</b>	
<b>Facility Move Details</b>	Health District Approval	Enter the name of the Health District representative who has approved the move to the new location, and vaccine storage and transportation arrangements.
	Can CCA be transferred from the old site to the new site?	Y <input type="checkbox"/> N <input type="checkbox"/> Tick <b>yes</b> if CCA has been confirmed to be transferred to the new location. Complete this section. Tick <b>no</b> if CCA has not been confirmed to be transferred to the new location and requires reassessment. Complete a site closure and new site set-up form.
	Address of new site	
	Is CCA expiry date current in CIR?	Y <input type="checkbox"/> N <input type="checkbox"/> If no, please update CIR.
	Is back up fridge current in CIR?	Y <input type="checkbox"/> N <input type="checkbox"/> If no, please update CIR.

# Appendix C:

## Facility/site closure

This information must be provided to NPHS Te Whatu Ora in the event of a facility or site choosing to no longer administer and distribute certain vaccine types.

Please take care and provide detail when completing the form below. Upon completion, please email this form to NPHS Te Whatu Ora service desk at: **help@c-19imms.min.health.nz**

For **user decommissioning**, please liaise with your Health District/Provider workforce lead or alternatively you can request the user decommissioning document from the NPHS Te Whatu Ora service desk.

### **The following definitions apply specifically to this form**

- **Vaccination facility**  
Where vaccines are shipped, stored, and distributed to sites.
- **Vaccination site**  
Where vaccines are administered.
- **CIR Suite**  
Includes all vaccination recording tools managed by NPHS Te Whatu Ora, the suite covers CIR, CIR Logistics portal, NIBS & Payments





# Facility and Site closure form

<b>Health District/Provider name</b> Please state the Health District/Provider the vaccination facility/site is attached to		
<b>A</b>	<b>Site closure</b>	
<b>Site</b>	1 Site name	
	2 Site address	
	3 Closure date	
	4 Reason for closure	
<b>B</b>	<b>Facility closure (if applicable)</b>	
<b>Facility</b>	5 Facility name	
	6 Facility address	
	7 Facility ID (if known)	
	8 Closure date	
	9 Reason for closure	
<b>C</b>	<b>Tick to confirm the closure of:</b>	
	Site: <input type="checkbox"/>	Facility: <input type="checkbox"/>
		Both: <input type="checkbox"/>
<b>D</b>	<b>If your site offers COVID and Flu vaccinations, tick to confirm the vaccine type you will no longer offer:</b>	
	COVID: <input type="checkbox"/>	Flu: <input type="checkbox"/>
		Both: <input type="checkbox"/>
<b>Return of excess stock</b>		
<p>1. Please conduct a stocktake of all assets relating to the COVID-19 Vaccination Rollout upon Facility/Site closure. Please send copies of this form to NPHS Te Whatu Ora Service Desk and your Health District Lead.</p> <p>2. The Health District lead should arrange a transfer of any remaining assets from the site which is closing to another site and capture this through raising a transfer order in CIR Inventory.</p> <p>3. Once there is zero stock on hand visible in CIR Inventory, the Health District logistics Lead should notify NPHS Te Whatu Ora to change the site status in CIR from Active to Closed.</p> <p><b>Note:</b> Once closed the site will not be accessible by CIR inventory users in the system.</p> <p>Providers must adhere to guidance provided in National Standards for Vaccine Storage and Transportation Providers 2017 and the 2021 Addendum when closing down a vaccine site/facility. Please refer to the links below for a copy.</p>		
Please tick to confirm these guidelines have been adhered to		Y <input type="checkbox"/>
		Please tick to confirm
<p>Once submitted, the site will be removed from all vaccination recording tools within the CIR suite. The site will no longer be visible through the logistics portal and if the site operates under a PPD model, they will be paid within the final cycle then removed from the contract.</p> <p>By completing this document, you agree that the Facility/Site will no longer be administering and/or ordering vaccines.</p>		

# Appendix D: Logistics and inventory management

NPHS Te Whatu Ora will maintain the COVID-19 Vaccination Immunisation Register (CIR) logistics module to support ongoing monitoring of inventory and demand. The image below shows the current process for distributing the vaccine to vaccination sites.

**Figure D.1 – vaccine distribution process**



Various vaccines have different storage conditions at each step of the process, see the Cold Chain Storage and shelf life in Section.

<b>Roles and Responsibilities</b>	<p><b>Vaccine Manufacturer</b> will ship trays to NZ's vaccine distributor, confirm temperature, then transfer ownership.</p>	<p><b>The Vaccine Distributor</b> will store at the optimal temperature for long term storage.</p>	<p><b>The Vaccine Distributor</b> will pick and pack and arrange transport to the vaccine facility for storage at +2°C to +8°C.</p>	<p><b>Sites</b> will forecast their daily volumes on a rolling weekly basis.</p>
	<p><b>NPHS Te Whatu Ora</b> will own the supply from here.</p>	<p><b>Health Districts</b> will maintain a demand plan for the upcoming four weeks and keep it up to date weekly.</p>	<p><b>Facilities</b> will receive and store vials at +2°C to +8°C in certified cold chain for later distribution to sites without cold chain.</p>	<p><b>Health Districts or Providers</b> may transport vials from their facilities to vaccination sites (within transportation time limits on vaccines if applicable).</p>
	<p><b>The Vaccine Distributor</b> will confirm the vaccine is undamaged and transfer to inventory management.</p>	<p><b>NPHS Te Whatu Ora</b> will confirm the order with the distributor to pack and transport to each delivery site.</p>	<p><b>Sites</b> may also receive and store vials at +2°C to +8°C in certified cold chain.</p>	

# Appendix E:

## NIP logistics overview/cheat sheets

### Regulations

- **COVID-19 vaccine ownership**  
All COVID-19 vaccine stock is owned by Pharmac.
- **Pharmacy licence**  
This allows Health District hospital pharmacies to pack down 10 vial cartons of COVID-19 vaccines into smaller quantities, but only for vaccination sites run by the Health District legal entity; that is, Health District hospital pharmacies can only pack down into smaller pack sizes for vaccination sites run by Health District employees.
- **Wholesale Licence**  
This allows Health District hospital pharmacies to supply COVID-19 vaccine by wholesale, in full cartons to non-health District vaccination sites outside their Health District legal entity. For the purposes of this, the definition of Health District means the Health District legal entity, not the geographical Health District boundary.

### Cold chain standards

- The **National Standards for Vaccine Storage and Transportation for Immunisation Providers 2017**, describes the standards and requirements for providers. The integrity of the cold chain is dependent upon:
  - the people who maintain and monitor the cold chain
  - the systems and processes used
  - and the equipment in which the vaccines are stored.
- **Cold chain accreditation**  
All immunisation providers are required to achieve accreditation (or Cold Chain Compliance, where applicable) if they need to store vaccine overnight. Assessors use this tool to ensure providers' cold chain practices and processes meet the required standards. See the **National Standards** for full details.
- An **Addendum** for ultra-cold vaccine storage of COVID-19 vaccine stock has been developed. Cold Chain Accreditation as per the addendum must be met before vaccines can be received.
- **Cold chain review group**  
Where a Health District hospital pharmacy needs to function outside of the boundaries of the national standards for cold chain storage and transportation, they can request NPHS Te Whatu Ora cold chain review group to be convened under urgency for advice. Any advice provided by this group will always be safe and ensure no compromise to the cold chain.

## Vaccine ordering

- **Registering new site/facility**

All sites/facilities need to be registered at least five days prior to the first required vaccine delivery. It is recommended the first delivery is used as a 'wet run' to vaccinate the vaccinators and to validate the delivery processes.

- **Order deadline**

Vaccine orders must be submitted before 10am the day before their allocated delivery day(s). Orders must be submitted in the CIR portal. The Health District lead needs to submit any urgent orders that are required prior to the next designated delivery day, as an 'out-of-cycle' delivery request to the CST Logistics Desk.

**Note:** If Health Districts need to check/QA vaccine site orders, ensure there is sufficient time for this process to be completed by 10am.

- **Consumable packs (being discontinued in march)**

Consumable packs containing needles and syringes can be ordered (100 kits) in addition to 'order as required' items.

## Vaccine handling

- **Receiving/sending at 2°C to 8°C**

COVID-19 vaccine arrives in validated cold-chain shipper boxes with a datalogger.

- **Shelf life**

See summary below, [and table 8.1](#) for full details.

- **Redistribution/transfers**

Vaccine stock is not to be redistributed between facilities and sites, unless requested by NPHS Te Whatu Ora or Health District Hospital pharmacy. Note: only HCL, DHL, and Health District hospital pharmacies have wholesale licences to support distribution of vaccine stock.

## Vaccine handling

- **Cold Chain accreditation and transportation**

All facilities must have a current cold chain accreditation and the expiry date recorded in the CIR. Providers must use temperature-monitored chilly bins to transport vaccines. A hard walled/robust chilly bin must be used for off-site clinics. For each chilly bin, monitor the temperature using either a digital minimum/maximum thermometer with an audible alarm, or a datalogger with a probe and external display. It must be possible to read the temperature without opening the chilly bin. All chilly bins and temperature monitors must be validated. Full details can be found in **section 7.3 of the national standards**.

- **Dataloggers**

Use a datalogger with a probe, external display and alarm to monitor the temperature of the vaccines throughout the time they are stored in a chilly bin. Set the datalogger to record the temperature every five minutes, and download, review and save the data after returning to the clinic. Full details can be found in **section 7.3 of the national standards**.

# Appendix F:

## Links to the National Immunisation Booking System

### COVID-19 Immunisation Register (CIR)

- All CIR training material can be found at  
[https://circlassrm-ncts.cs116.force.com/cir/s/recordlist/Knowledge\\_\\_kav/00B5O000001CNbyUAG](https://circlassrm-ncts.cs116.force.com/cir/s/recordlist/Knowledge__kav/00B5O000001CNbyUAG)

### Individual guides

#### **NIBS**

- <https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Bookings>
- <https://circlassrm-ncts.cs116.force.com/cir/s/article/CIR-Bookings-Not-NHI-Matched-Quick-Step-Guide>

#### **Accenture Vaccine Management System (AVMS)**

- <https://circlassrm-ncts.cs116.force.com/cir/s/article/NIBS-Site-Admin-Managing-Overrides-exceptions-Guide>
- <https://circlassrm-ncts.cs116.force.com/cir/s/article/NIBS-Site-Admin-Managing-Capacity-Guide>

#### **Other information**

For any information which is not included in these documents, the Health District is advised to communicate with NPHS Te Whatu Ora.

This guide will be amended as required and the latest version will be made available via:

- [https://circlassrm-ncts.cs116.force.com/cir/s/recordlist/Knowledge\\_\\_kav/00B5O000001CNbyUAG](https://circlassrm-ncts.cs116.force.com/cir/s/recordlist/Knowledge__kav/00B5O000001CNbyUAG)

# Appendix G:

## Vaccination site screening questions

We encourage you to screen both staff and consumers for risk of exposure to COVID-19 and COVID-19 symptoms. Screening is critical to breaking the chain of transmission of COVID-19 and maintaining staff and consumer safety.

**Figure G.1** below details the recommended screening questions and process to create a lower risk environment for transmission of COVID-19 and to ensure PPE advice is appropriate.

**Please note:**

- In the event of COVID-19 Alert Level changes, additional advice will be formulated by local public health units and NPHS Te Whatu Ora.
- Any consumer with a confirmed COVID-19 infection should not be vaccinated until they have had the appropriate recovery period (see Immunisation Handbook or consult with IMAC).
- Any consumer that has answered 'yes' to the screening questions below, is considered high risk for the transmission of COVID-19 and deferral is recommended.
- If a provider wishes to vaccinate a higher risk consumer (someone who answered yes below), providers should follow the 'vaccination in high-risk or screen 'positive' consumers' section in the Operating Guidelines.

### Figure G.1– recommended screening questions

#### Q1 – Do you have symptoms of COVID-19?

Follow link to **COVID-19 Case definition**

If a client has any symptoms suggestive of COVID-19, defer vaccination and do not permit entry to the site. Advise them to follow recommendations and guidance from NPHS Te Whatu Ora/public health services. Recommend they get a test and self-isolate pending the result.



If no symptoms, continue to the next question.

#### Q2 – Do you live with someone who has COVID-19?

If an individual lives with someone who has COVID-19, they are considered a household contact do not permit entry to the site and advise them to follow recommendations and guidance from NPHS Te Whatu Ora/public health services.



If no symptoms, continue to the next question.

#### Q3 – Have you been requested to stay at home, to self-isolate or are under an isolation order?

If yes, defer vaccination and do not permit entry to the site. Recommend continuing to follow the stay at home/self-isolation plan.



If no symptoms, continue to the next question.

#### Q4 – Are you currently waiting on a COVID-19 test result?


If yes, defer vaccination and do not permit entry to the site. Recommend rebooking once a negative test result has been received, and they have been told they no longer need to stay at home/self-isolate.



If no, proceed to vaccinate as per the Operating Guidelines.

# Appendix H: Supported decision-making process

Figure H.1 – support to consent



# Appendix I:

## **NIP Adverse Event Process**

This Appendix includes

1. Introduction
2. Process Steps
3. Severity Assessment Code (SAC) examples
4. Provider with the NIP incident notification form.



# Provider and Programme Lead Clinicians

## Purpose

The National Immunisation Programme (NIP) implementation phase is based on a devolved service delivery model. The NIP Clinical Lead is committed to supporting a person-centred, safe and high-quality Programme with all Programme providers.

To support a provider when a serious adverse event occurs, the following process includes timely notification to the Programme and consideration of NIP support to the provider.

The following detail outlines the notification process and describes roles/responsibilities of NIP provider lead clinicians in relation to COVID-19 vaccination-related serious adverse event<sup>2</sup> or a serious adverse event following immunisation<sup>3</sup>.

## Scope

This process outlines the notification of NIP adverse events and uses severity assessment code (SAC) ratings. Any of the following must follow this notification process:

- SAC 1 (e.g., Anaphylaxis resulting in death or permanent loss of function)
- SAC 2 (e.g., Serious adverse reaction with delayed administration of adrenaline or delayed presence of emergency services)
- SAC 3 (e.g., Medication error, vaccine dilution error, or dose error)
- Several similar or close sequenced SAC 4 events (e.g., Breach of confidentiality).
- Near miss with likely significant consequences

**Note:** For more examples of the SAC ratings please refer to the table below.

This protocol aligns with existing expectations of health and disability service providers under the Health and Disability Services (Safety) Act 2001, as articulated by the Health Quality & Safety Commission, whereby those who voluntarily comply are expected to:

1. Report serious adverse events (SAC rating 1 and 2) and events on the Always Report and Review list to the Commission, using the adverse event brief – part A reporting form. This report should be made within 15 working days of notification of the event to the provider.
2. Undertake formal investigation of serious adverse events (SAC 1 and 2) and events on the Always Report and Review list and send review findings and recommendations to the Commission, using the adverse event brief – part B reporting form. This report should be made within 70 working days of notification of the event to the provider.

## Exclusions

This NIP serious adverse event process does not apply to other NIP non-clinical incident types e.g., equipment or vaccine damage/loss.

The notification process is not a substitute for the provider's responsibility concerning an adverse event including their normal processes of reporting, reviewing and open communication with the affected person. The outcome may recommend clinical and quality continuous improvement actions.

---

<sup>2</sup> An adverse event is an incident resulting in harm, or with the potential to result in harm to a health consumer.

<sup>3</sup> Adverse event following immunisation (AEFI) - an untoward medical event which follows immunisation and does not necessarily have a causal relationship with the administration of the vaccine. The adverse event may be an unfavourable or unintended sign, abnormal laboratory finding, symptom or disease

# Process Steps

Pharmacovigilance	Timeframe
Ensure COVID-19 CARM report is completed for any suspected AEFI. <a href="https://report.vaccine.covid19.govt.nz">CARM Resource</a> <a href="https://report.vaccine.covid19.govt.nz">https://report.vaccine.covid19.govt.nz</a>	Day 1 ( < 8 hours)
Participate in follow-up activities with CARM if required.	On contact by CARM



Notification to NPHS Te Whatu Ora NIP and provider leads	Timeframe
Commence reporting process. You should use the attached provider or organisation process steps and ensure you identify a <u>preliminary</u> SAC rating. <a href="#">Programme Resource: NIP SAC examples in table below</a> <a href="#">HQSC resource A Guide to the National Adverse Events Reporting Policy 2017</a>	Day 1 ( < 8 hours)
<b>Notify NIP Programme via email address:</b> <a href="mailto:NIP.incidentnotification@health.govt.nz">NIP.incidentnotification@health.govt.nz</a> <ul style="list-style-type: none"> <li>• <b>Attach the completed:</b> Provider with NIP incident notification form (sections A and B) <ul style="list-style-type: none"> <li>○ Email Subject: NIP Adverse Event Notification</li> </ul> </li> <li>• Please refer to the relevant incident toolkit which can be found on the Connex 'Mahi Tahī' platform or your provider's Clinical/Quality Lead.</li> </ul> <a href="#">Programme Resource: Provide with NIP incident notification form<sup>4</sup></a>	Expedited ( < 48 hours)



<b>Plan and execute open communication with affected consumer/s<sup>5</sup></b>	<b>Within 7 working days</b>
---	------------------------------



Investigation and reporting outcomes	Timeframe
<ul style="list-style-type: none"> <li>• Investigate the incident using the provider or organisation's clinical quality and safety governance process, and in accordance with HQSC expectations.</li> <li>• Inform NIP on investigation findings and recommendations.</li> <li>• This includes confirming the final SAC rating.</li> </ul> <a href="https://www.hqsc.govt.nz/our-work/system-safety/adverse-events/">HQSC resource</a> <a href="https://www.hqsc.govt.nz/our-work/system-safety/adverse-events/">https://www.hqsc.govt.nz/our-work/system-safety/adverse-events/</a>	Commenced ( < 24 hours)  Reporting to HQSC according to timeframes above.
<b>If required, please arrange an <a href="#">ACC treatment injury claim</a>. Also see the <a href="#">Treatment injury claim lodgement guide</a> and the <a href="#">Treatment Injury Flowchart</a>.</b>	
<b>Updating of NIP incident form and send an update to NPHS Te Whatu Ora NIP</b>	Ongoing/ until closed

<sup>4</sup> This is the notification form for all incident types including serious adverse events & AEFI.

<sup>5</sup> As a guide, the Health Quality and Safety Commission's "Root Cause Analysis for clinical incidents - A Practical Guide" have the expectation for communication with affected consumers during week 1 – 2 of the incident investigations.

**Provider please:**

As an adverse event, either following immunisation or other cause, please arrange for open communication with the affected person/s.

If required, please arrange ACC treatment injury claim per ACC2152 form: <https://www.acc.co.nz/assets/provider/3e3bd2aded/acc2152-treatment-injury-claim.doc>

SAC 1 Death or permanent severe loss of function	SAC 2 Permanent major or temporary severe loss of function
<ul style="list-style-type: none"> <li>• Medication or dose error resulting in death or causing renal failure and need for permanent renal replacement therapy</li> <li>• Anaphylaxis resulting in death or permanent loss of function</li> <li>• Wrong site of vaccine resulting in removal of healthy limb or organ</li> <li>• Delayed referral, treatment resulting in treatment options limited to palliation (delay direct contributor)</li> <li>• Delayed recognition of patient deterioration resulting in permanent disability or death</li> </ul>	<ul style="list-style-type: none"> <li>• Fall resulting in fracture</li> <li>• Serious adverse reaction with delayed administration of adrenaline or delayed presence of emergency services</li> <li>• Delayed recognition of patient deterioration resulting in unplanned transfer to intensive care or to another hospital for higher acuity care, cardiopulmonary resuscitation and/or intubation</li> <li>• Medication or vaccine dose error resulting in major harm (e.g., requiring dialysis, intervention to sustain life, anaphylaxis)</li> <li>• Consumer serious assault occurring within vaccination care setting when a known safety plan is not upheld (e.g., protection order)</li> <li>• A vaccination incident affecting &gt; 1 consumer</li> </ul>
SAC 3 Permanent moderate or temporary major loss of function	SAC 4 Requiring increased level of care OR no injury, no increased level of care; includes near misses
<ul style="list-style-type: none"> <li>• Fall resulting in laceration requiring sutures</li> <li>• Failure of essential service with moderate consequence to consumer</li> <li>• Medication error, vaccine dilution error, or dose error</li> <li>• Temporary nerve damage or pain from vaccine administration</li> <li>• Severe injection site infection</li> <li>• Vasovagal event following immunisation resulting in injury</li> <li>• Never events: wrong vaccine, early vaccination doses &amp; underage vaccination</li> </ul>	<ul style="list-style-type: none"> <li>• Additional monitoring, investigations, or interventions due to the event post vaccination</li> <li>• Medication, vaccine dilution or dose error resulting in no increased level of care or monitoring- not reaching the consumer is a near miss</li> <li>• Breach of confidentiality</li> <li>• Near miss events</li> </ul>

Version 4: Adapted for the National Immunisation Programme (NIP) from Severity Assessment Code (SAC) examples 2019–20 | Health Quality & Safety Commission 2019. This list is guidance only.

## NIP incident notification form

<b>Notify and attach this completed form to:</b> nip.incidentnotification@health.govt.nz			
<b>Email Subject:</b> NIP Adverse Event Notification			
<b>Verified from the NIP Detecting Failsafe Report:</b> Y <input type="checkbox"/> N <input type="checkbox"/>			
<b>Section A – Provider notification details</b>			
Provider or Health District to complete information below			
Incident date/ time			
Date/ time reported			
Site		Health District	
<b>Person reporting incident:</b>			
Name			
Contact phone number/s			
Email address			
<b>Section B – Description (Provider to complete)</b>			
Type of incident / adverse event / AEFI (it's possible two of the four options apply)			
Near miss <input type="checkbox"/>	Incident <input type="checkbox"/>	Serious adverse event <input type="checkbox"/>	AEFI <input type="checkbox"/>
Vaccine type and dose (e.g., Comirnaty 30mcg grey cap):	Dose details (circle):	Primary Dose 1 / Dose 2 / Dose 3 Booster 1 / Booster 2 Other:	
Age of consumer:		Ethnicity:	
Have the Health District's/Provider's Clinical Lead or Quality Lead been notified?			Y <input type="checkbox"/> N <input type="checkbox"/>
If there is an adverse event following immunisation or a medication error, has this been reported to CARM?			Y <input type="checkbox"/> N <input type="checkbox"/>
Has IMAC been contacted for advice and given to the consumer:			Y <input type="checkbox"/> N <input type="checkbox"/>
Has CIR been entered correctly to reflect actual dose given?			Y <input type="checkbox"/> N <input type="checkbox"/>
Has a preliminary investigation been undertaken? List details below			Y <input type="checkbox"/> N <input type="checkbox"/>
Has the consumer been informed and received an apology?			Y <input type="checkbox"/> N <input type="checkbox"/>
Assign a preliminary SAC rating (circle one):			SAC 1 / 2 / 3 / 4
<ul style="list-style-type: none"> <li><b>Incident</b> means any unplanned event resulting in, or having a potential for injury, ill health, damage or other loss, an incident includes an accident.</li> <li><b>Adverse event</b> is an incident resulting in harm, or with the potential to result in harm to a health consumer. Please assign an adverse event SAC rating. Report a SAC 1,2 or 3 SAC event, a cluster of SAC 3/ 4 events +/- near misses.</li> </ul>			

- **Adverse event following immunisation (AEFI)** is an untoward medical event which follows immunisation and does not necessarily have a causal relationship with the administration of the vaccine. The adverse event may be an unfavourable or unintended sign, abnormal laboratory finding, symptom or disease.

**Provider please note:**

- Include information regarding open communication with an affected consumer, including date completed
- Include your findings in the actions you have taken to prevent reoccurrence
- Update this section of the form over time as incident investigation is progressed and then closed

**Please provide as much detail of the incident as possible:**

What went wrong? Were there any contributing factors? What were the immediate actions taken? What advice were you given and from whom? What changes will you be making to prevent this happening again? What follow up has been arranged for the consumer?

If the consumer received an early dose, please provide the number of days between doses.

**Reviewed by (name and role):**

Clinical Lead or Quality Lead

# Appendix J:

## Risk mitigations for vaccination sites

**Table L1 – risk mitigations**

Actions Required at all levels	Supporting Document
<ul style="list-style-type: none"> <li>Adapt processes as required for screening of staff, consumers, and support people to capture COVID-19 symptoms, travel history, and/or attendance at locations of interest, if they have been directed to have a test or are awaiting a test result. Redirect symptomatic consumers or those with contact history for testing in line with Ministry of Health guidance.</li> </ul>	<ul style="list-style-type: none"> <li><b>Operating Guidelines</b> Refer to the <b>Vaccination Site Screening Questions</b> section above.</li> </ul>
<ul style="list-style-type: none"> <li>Robust communication strategy to regularly inform staff and consumers of Programme and service delivery changes.</li> </ul>	<ul style="list-style-type: none"> <li><b>COVID-19: Q&amp;A for primary health care workers.</b></li> </ul>
<ul style="list-style-type: none"> <li>Promote staff awareness of resources to maintain up-to-date knowledge of national COVID-19 related information.</li> </ul>	<ul style="list-style-type: none"> <li><b>Āwhina App</b></li> </ul>
<ul style="list-style-type: none"> <li>Oversee and manage safe access to the site and queue management.</li> </ul>	<ul style="list-style-type: none"> <li><b>Operating Guidelines</b></li> </ul>
<ul style="list-style-type: none"> <li>Orientation and Adherence to Infection Prevention and Control (IPC) guidance, including hand hygiene, and Personal Protective Equipment (PPE) guidelines for various situations. These must be available and understood.</li> </ul>	<ul style="list-style-type: none"> <li><b>Five Moments of Hand Hygiene</b></li> <li><b>FAQ regarding IPC and PPE</b></li> <li><b>PPE use in Health and Disability Care Settings</b></li> </ul>
<ul style="list-style-type: none"> <li>Plans to support adequate and safe staffing to deliver services.</li> </ul>	<ul style="list-style-type: none"> <li><b>Operating Guidelines</b></li> </ul>
<ul style="list-style-type: none"> <li>Ensure there is sufficient internet connectivity to enable use of the CIR and other technology in all relevant areas of the site. It may be necessary to use mobile Wi-Fi hotspots.</li> </ul>	<ul style="list-style-type: none"> <li><b>Operating Guidelines</b></li> </ul>
<ul style="list-style-type: none"> <li>Staff wellness: Staff must be discouraged from attending work when unwell and must be encouraged to be up to date with occupationally relevant vaccinations.</li> </ul>	
<ul style="list-style-type: none"> <li>Ensure that environmental safety considerations, including ventilation, are adequately appraised.</li> </ul>	

# Document version control

## Revision History

Version	Date	Section/ Appendix	Summary of Changes
48.0	9/12/22	<b>Section A</b>	
		9.2	Updated information and photos to reflect the upgraded temperate monitoring device which will commence being used in DHL deliveries from 28 November 2022.
		<b>No changes to sections B, C, D, E or appendices</b>	
49.0	11/01/23	<b>Section A</b>	
		<b>Roles and responsibilities</b>	Changed purchasing vaccines from Pfizer to purchasing vaccines
		<b>Table 7.3</b>	Added Nuvaxovid and Pfizer Paediatric to the Instructions for the Preparation and Administration of Vaccines and updated links to the IMAC resources
		<b>9.2.2</b>	Added Nuvaxovid and Pfizer Paediatric to the Instructions for the Preparation and Administration of Vaccines and updated links to the IMAC resources
		<b>No changes to sections B, C, D, E or appendices</b>	
50.0	08/02/23	7.1	Removed COVID19 Tracer app QR code section
		Table 9.5	Updated Econolog temperature reading device photos to show different colour variations
		<b>Section B</b>	
		B	Section B has been merged with sections C (paediatric Pfizer) and D (Novavax), Comirnaty maroon cap vaccine operational information has been added (pages 81-84).
		21	COVID-19 vaccine pathway to vaccination section has been refreshed, there is now only one section for this instead of one for each vaccine.
		21.2	Additional sentence added for consumers under observation following concomitant vaccination.
		<b>Section C</b>	
		29.5	Updated to reflect Comirnaty maroon cap rollout
		<b>Appendices</b>	
		Appendix K	Removed as no longer required